Fields Institute Communications
Volume **00**, 0000

arXiv:1212.4308v1 [math.NT] 18 Dec 2012

Modularity of Calabi–Yau varieties: 2011 and beyond

Noriko YUI
Department of Mathematics and Statistics
Queen's University
Kingston, Ontario, Canada K7L 3N6
E-mail: yui@mast.queensu.ca

Abstract. This paper presents the current status on modularity of Calabi–Yau varieties since the last update in 2003. We will focus on Calabi–Yau varieties of dimension at most three. Here modularity refers to at least two different types: arithmetic modularity and geometric modularity. These will include: (1) the modularity (automorphy) of Galois representations of Calabi–Yau varieties (or motives) defined over $\mathbb{Q}$ or number fields, (2) the modularity of solutions of Picard–Fuchs differential equations of families of Calabi–Yau varieties, and mirror maps (mirror moonshine), (3) the modularity of generating functions of invariants counting certain quantities on Calabi–Yau varieties, and (4) the modularity of moduli for families of Calabi–Yau varieties. The topic (4) is commonly known as geometric modularity.

Discussions in this paper are centered around arithmetic modularity, namely on (1), and (2), with a brief excursion to (3).

Introduction

These are notes of my introductory lectures at the Fields workshop on "Arithmetic and Geometry of K3 surfaces and Calabi–Yau threefolds", at the Fields Institute from August 16 to August 25, 2011. My goal is two-fold:

- Present an update on the recent developments on the various kinds of modularity associated to Calabi–Yau varieties. Here "recent" developments means various developments since my article [65] published in 2003.
- Formulate conjectures and identify future problems on modularity and related topics.

1991 *Mathematics Subject Classification.* 11G42, 11F80, 11G40, 14J15, 14J32, 14J33.

Key words and phrases. Elliptic curves, K3 surfaces, Calabi–Yau threefolds, CM type Calabi–Yau varieties, Galois representations, modular (cusp) forms, automorphic inductions, geometry and arithmetic of moduli spaces, Hilbert and Siegel modular forms, families of Calabi–Yau varieties, mirror symmetry, mirror maps, Picard–Fuchs differential equations.

The author was supported in part by NSERC Discovery Grant.

©0000 American Mathematical Society

My hope is to motivate young (as well as mature) researchers to work in this fascinating area at the interface of arithmetic, geometry and physics around Calabi–Yau varieties.

0.1 Brief history since 2003. The results and discoveries in the last ten years on Calabi–Yau varieties, which will be touched upon in my lectures, are listed below.

- The modularity of the 2-dimensional Galois representations associated to Calabi–Yau varieties defined over $\mathbb{Q}$.
- The modularity of highly reducible Galois representations associated to Calabi–Yau threefolds over $\mathbb{Q}$.
- The automorphy of higher dimensional Galois representations arising from CM type Calabi–Yau varieties (automorphic induction).
- Appearance of various types of modular forms in Mirror Symmetry as generating functions counting some mathematical/physical quantities.
- Modularity of families of Calabi–Yau varieties (solutions of Picard–Fuchs differential equations, monodromy groups, mirror maps).
- Moduli spaces of Calabi–Yau families, and higher dimensional modular forms (e.g., Siegel modular forms).

0.2 Plan of lectures. Obviously, due to time constraints, I will not be able to cover all these topics in my two introductory lectures. Thus, for my lectures, I plan to focus on recent results on the first three (somewhat intertwined) items listed above, and with possibly very brief interludes to the rest if time permits;

This article includes my two lectures delivered at the workshop, as well as some subjects/topics which I was not able to cover in my lectures. However, I must emphasize that this note will not touch upon geometric modularity.

0.3 Disclaimer. Here I will make the disclaimer that the topics listed above are by no means exhaustive; it may be the case that I forget to mention some results, or not give proper attributions. I apologize for these oversights.

0.4 Calabi–Yau varieties: Definition.

Definition 0.1 Let X be a smooth projective variety of dimension d defined over $\mathbb{C}$. We say that X is a *Calabi–Yau* variety if

- $H^i(X,\mathcal{O}_X)=0$ for every i, $0<i<d$.
- The canonical bundle K_X is trivial.

We introduce the *Hodge numbers* of X:

$$h^{i,j}(X):=\dim_{\mathbb{C}}H^j(X,\Omega^i_X),\ 0\leq i,j\leq d.$$

Then we may characterize a Calabi–Yau variety of dimension d in terms of its Hodge numbers.

A smooth projective variety X of dimension d over $\mathbb{C}$ is called a *Calabi–Yau* variety if

- $h^{i,0}(X)=0$ for every i, $0<i<d$.
- $K_X\simeq\mathcal{O}_X$, so that the geometric genus of X,

$$p_g(X):=h^{0,d}(X)=\dim_{\mathbb{C}}H^0(X,K_X)=\dim_{\mathbb{C}}H^0(X,\mathcal{O}_X)=1.$$

Numerical characters of Calabi–Yau varieties of dimension d

- Betti numbers: For i, $0 \leq i \leq 2d$, the i-th Betti number of X is defined by
$$B_i(X) := \dim_{\mathbb{C}} H^i(X,\mathbb{C}).$$
There is Poincaré duality for $H^i(X,\mathbb{C})$; that is,
$$H^i(X,\mathbb{C}) \times H^{2d-i}(X,\mathbb{C}) \to \mathbb{C} \quad \text{for every } i,\ 0 \leq i \leq d$$
is a perfect pairing. This implies that
$$B_i(X) = B_{2d-i}(X), \text{for } i,\ 0 \leq i \leq d$$
- Hodge numbers: They are defined in Definition 0.1 above. There is the symmetry of Hodge numbers: For $0 \leq i, j \leq d$,
$$h^{i,j}(X) = h^{j,i}(X) \quad \text{by complex conjugation,}$$
and
$$h^{i,j}(X) = h^{d-i,d-j}(X) \quad \text{by Serre duality.}$$
- There is a relation among Betti numbers and Hodge numbers, as a consequence of the Hodge decomposition:
$$B_k(X) = \sum_{i+j=k} h^{i,j}(X).$$
- The Euler characteristic of X is defined by
$$E(X) := \sum_{k=0}^{2d} (-1)^k B_k(X).$$

Example 0.2 (a) Let $d = 1$. The first condition is vacuous. The second condition says that $p_g(X) = 1$. So dimension 1 Calabi–Yau varieties are elliptic curves. The Hodge diamond of elliptic curves is rather simple.

$$\begin{array}{ccc} & 1 & \\ 1 & & 1 \\ & 1 & \end{array} \qquad \begin{array}{l} B_0(X) = 1 \\ B_1(X) = 2 \\ B_2(X) = 1 \end{array}$$

The Euler characteristic of X is given by
$$E(X) = B_0(X) - B_1(X) + B_2(X) = 0.$$

(b) Let $d = 2$. The first condition is $h^{1,0}(X) = 0$ and the second condition says that $h^{0,2}(X) = p_g(X) = 1$. So dimension 2 Calabi–Yau varieties are K3 surfaces. The Hodge diamond of K3 surfaces is of the form:

$$\begin{array}{ccccc} & & 1 & & \\ & 0 & & 0 & \\ 1 & & 20 & & 1 \\ & 0 & & 0 & \\ & & 1 & & \end{array} \qquad \begin{array}{l} B_0(X) = 1 \\ B_1(X) = 0 \\ B_2(X) = 20 \\ B_3(X) = 0 \\ B_4(X) = 1 \end{array}$$

The Euler characteristic of X is given by
$$E(X) = \sum_{k=0}^{4} (-1)^k B_k(X) = 1 + 22 + 1 = 24.$$

(c) Let $d = 3$. The first condition says that $h^{1,0}(X) = h^{2,0}(X) = 0$ and the second condition implies that $h^{0,3}(X) = p_g(X) = 1$. The Hodge diamond of X is given as follows:

$$\begin{array}{ccccccc}
 & & & 1 & & & \\
 & & 0 & & 0 & & \\
 & 0 & & h^{1,1}(X) & & 0 & \\
1 & & h^{2,1}(X) & & h^{1,2}(X) & & 1 \\
 & 0 & & h^{2,2}(X) & & 0 & \\
 & & 0 & & 0 & & \\
 & & & 1 & & &
\end{array}$$

$$\begin{aligned}
B_0(X) &= 1\\
B_1(X) &= 0\\
B_2(X) &= h^{1,1}(X)\\
B_3(X) &= 2(1+h^{2,1}(X))\\
B_4(X) &= h^{2,2}(X) = h^{1,1}(X)\\
B_5(X) &= 0\\
B_6(X) &= 1
\end{aligned}$$

The Euler characteristic is given by

$$E(X) = \sum_{k=0}^{6}(-1)^k B_k(X) = 2(h^{1,1}(X) - h^{2,1}(X)).$$

It is not known if there exist absolute constants that bound $h^{1,1}(X)$, $h^{2,1}(X)$, and hence $|E(X)|$. The currently known bound for $|E(X)|$ is 960.

(d) Here are some typical examples of families of Calabi–Yau varieties defined by hypersurfaces.

d	CY variety of dim d	CY varieties of dim d
1	$X_0^3+X_1^3+X_2^3+3\lambda X_0X_1X_2$	$y^2=f_3(x)$
2	$X_0^4+X_1^4+X_2^4+X_3^4+4\lambda X_0X_1X_2X_3$	$z^2=f_6(x,y)$
3	$X_0^5+X_1^5+X_2^5+X_3^5+X_4^5+5\lambda X_0X_1X_2X_3X_4$	$w^2=f_8(x,y,z)$

The equations in the second column are generic polynomials in projective coordinates; while those in the third column are in affine coordinates and the f_i are smooth polynomials of degree i in affine coordinates.

We have a vast source of examples of Calabi–Yau threefolds via toric construction and other methods. The upper record for the absolute value of the Euler characteristic of all these Calabi–Yau threefolds is 960, though there is neither reason nor explanation for this phenomenon.

1 The modularity of Galois representations of Calabi–Yau varieties (or motives) over $\mathbb{Q}$.

We now consider smooth projective varieties defined over $\mathbb{Q}$ (say, by hypersurfaces, or complete intersections). We say that $X/\mathbb{Q}$ is a Calabi–Yau variety of dimension d over $\mathbb{Q}$, if $X\otimes_{\mathbb{Q}}\mathbb{C}$ is a Calabi–Yau variety of dimension d. A Calabi–Yau variety X over $\mathbb{Q}$ has a model defined over $\mathbb{Z}[\frac{1}{m}]$ (with some $m\in\mathbb{N}$), and this allows us to consider its reduction modulo primes. Pick a prime p such that $(p,m)=1$, and define the reduction of X modulo p, denoted by $X \mod p$. We say that p is a *good* prime if $X \mod p$ is smooth over $\bar{\mathbb{F}}_p$, otherwise p is *bad*. For a good prime p, let Fr_p denote the Frobenius morphism on X induced from the p-th power map $x\mapsto x^p$.

Let ℓ be a prime different from p. Then, for each i, $0\le i\le 2d$, Fr_p induces an endomorphism Fr_p^* on the i-th ℓ-adic étale cohomology group $H^i_{et}(\bar{X}_p,\mathbb{Q}_\ell)$, where $\bar{X}_p := X\otimes_{\mathbb{F}_p}\bar{\mathbb{F}}_p$. Grothendieck's specialization theorem gives an isomorphism $H^i_{et}(\bar{X}_p,\mathbb{Q}_\ell)\cong H^i_{et}(\bar{X},\mathbb{Q}_\ell)$, where $\bar{X}:=X\otimes_{\mathbb{Q}}\bar{\mathbb{Q}}$. Then the comparison theorem gives $H^i_{et}(\bar{X},\mathbb{Q}_\ell)\otimes_{\mathbb{Q}_\ell}\mathbb{C}\cong H^i(X\otimes_{\mathbb{Q}}\mathbb{C},\mathbb{C})$ so that $\dim_{\mathbb{Q}_\ell} H^i_{et}(\bar{X},\mathbb{Q}_\ell)=B_i(X)$. There is Poincaré duality for $H^i_{et}(\bar{X},\mathbb{Q}_\ell)$, that is,

$$H^i_{et}(\bar{X},\mathbb{Q}_\ell)\times H^{2d-i}_{et}(\bar{X},\mathbb{Q}_\ell)\to\mathbb{Q}_\ell \quad \text{for every } i,\ 0\le i\le 2d$$

is a perfect pairing. Let

$$P^i_p(T):=\det(1-\mathrm{Fr}^*_p T\mid H^i_{et}(\bar{X},\mathbb{Q}_\ell))$$

be the reciprocal characteristic polynomial of Fr^*_p. (Here T is an indeterminate.) Then the Weil Conjecture (Theorem) asserts that

- $P^i_p(T)\in 1+T\mathbb{Z}[T]$. Moreover, $P^i_p(T)$ does not depend on the choice of ℓ.
- $P^i_p(T)$ has degree $B_i(X)$.
- $P^{2d-i}_p(T)=\pm P^i_p(p^{d-\frac{i}{2}}T)$ for every i, $0\le i\le d$.
- If we write
$$P^i_p(T)=\prod_{j=1}^{B_i}(1-\alpha_{ij}\,T)\in\bar{\mathbb{Q}}[T]$$
then α_{ij} are algebraic integers with $|\alpha_{ij}|=p^{i/2}$ for every i, $0\le i\le 2d$.

Now we will bring in the absolute Galois group $G_{\mathbb{Q}}:=\mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. There is a continuous system of ℓ-adic Galois representations

$$\rho^i_{X,\ell}:G_{\mathbb{Q}}\to GL(H^i_{et}(\bar{X},\mathbb{Q}_\ell))$$

sending the (geometric) Frobenius Fr_p^{-1} to $\rho^i(\mathrm{Fr}_p^{-1})$. The (geometric) Frobenius $\rho^i(\mathrm{Fr}_p^{-1})$ has the same action as the Frobenius morphism Fr^*_p on the étale cohomology $H^i_{et}(\bar{Z},\mathbb{Q}_\ell)$. We define its L-series $L(\rho^i_{X,\ell},s):=L(H^i_{et}(\bar{X},\mathbb{Q}_\ell),s)$ for each i, $0\le i\le 2d$, where s is a complex variable.

We will now define the L-series of X.

Definition 1.1 The i-th (cohomological) L-series of X is defined by the Euler product

$$\begin{aligned}L_i(X,s)&:=L(H^i_{et}(\bar{X},\mathbb{Q}_\ell),s)\\ &:=(*)\prod_{p:p\neq\ell}P^i_p(p^{-s})^{-1}\times(\text{factor corresponding to } \ell=p)\end{aligned}$$

where the product is taken over all good primes different from ℓ and $(*)$ corresponds to factors of bad primes. For $\ell=p$, we may choose another good prime $\ell\neq p$, or we can use some p-adic cohomology (e.g., crystalline cohomology) to define the factor.

For $i=d$, we write simply $L(X,s)$ for $L_d(X,s)$ if there is no danger of ambiguity.

Remark 1.2 We may define (for a good prime p) the zeta-function $\zeta(X_p,T)$, of a Calabi–Yau variety X_p defined over $\mathbb{F}_p$ by counting the number of rational points on all extensions of $\mathbb{F}_p$:

$$\zeta(X_p,T):=\exp\left(\sum_{n=1}^{\infty}\frac{\#X_p(\mathbb{F}_{p^n})}{n}T^n\right)\in\mathbb{Q}(T).$$

Then by Weil's conjecture, it has the form:

$$\zeta(X_p,T)=\frac{P_1(T)P_3(T)\cdots P_{2d-1}(T)}{P_0(T)P_2(T)\cdots P_{2d}(T)},$$

where we put (to ease the notation) $P_i(T) = P_p^i(T)$ for $i = 1, \ldots, 2d$.

Let X be a Calabi–Yau variety of dimension d defined over $\mathbb{Q}$. For a good prime p, Fr_p acts on $X \mod p$, and it will induce a morphism Fr_p^* on $H^i_{et}(\bar{X}_p, \mathbb{Q}_\ell) \simeq H^i_{et}(\bar{X}, \mathbb{Q}_\ell)$. Define the trace $t_i(p)$ by

$$t_i(p) := \mathrm{Trace}(\mathrm{Fr}_p^* \,|\, H^i_{et}(\bar{X}, \mathbb{Q}_\ell)), \quad \text{for } i,\ 0 \le i \le 2d.$$

Then $t_i(p) \in \mathbb{Z}$ for every i, $0 \le i \le 2d$. The Lefschetz fixed point formula gives a relation between the number of $\mathbb{F}_p$-rational points on X and traces:

$$\# X(\mathbb{F}_p) = \sum_{i=0}^{2d} (-1)^i t_i(p).$$

Example 1.3 (a) Let $d = 1$ and let E be an elliptic curve defined over $\mathbb{Q}$. For a good prime p,

$$\# E(\mathbb{F}_p) = \sum_{i=0}^{2} (-1)^i t_i(p) = t_0(p) - t_1(p) + t_2(p) = 1 + p - t_1(p),$$

where

$$|t_1(p)| \le 2p^{1/2}.$$

Then we have

$$P_p^0(T) = 1 - T,\ P_p^2(T) = 1 - pT,\ P_p^1(T) = 1 - t_1(p)T + pT^2.$$

The L-series of E is then given by

$$\begin{aligned} L(E,s) &= (*) \prod_{p:good} P_p^1(p^{-s})^{-1} \\ &= (*) \prod_{p:good} \frac{1}{1 - t_1(p)p^{-s} + p^{1-2s}}. \end{aligned}$$

Expanding out, we may write

$$L(E,s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s} \quad \text{with } a_1 = 1 \text{ and } a(n) \in \mathbb{Z}.$$

So

$$a(p) = t_1(p) \quad \text{for every good prime } p.$$

(b) Let $d = 2$, and let X be a K3 surface defined over $\mathbb{Q}$. Let $NS(X)$ denote the Néron–Severi group of X generated by algebraic cycles. It is a free finitely generated abelian group, and

$$NS(X) = H^2(X, \mathbb{Z}) \cap H^{1,1}(X)$$

so that the rank of $NS(X)$ (called the Picard number of X), denoted by $\rho(X)$, is bounded above by 20. Let $T(X)$ be the orthogonal complement of $NS(X)$ in $H^2(X, \mathbb{Z})$ with respect to the intersection pairing. We call $T(X)$ the group of transcendental cycles of X. We have the decomposition

$$H^2(X, \mathbb{Z}) \otimes \mathbb{Q}_\ell = (NS(X) \otimes \mathbb{Q}_\ell) \oplus (T(X) \otimes \mathbb{Q}_\ell).$$

and this will enable us decompose the L-series of X as follows:

$$L(X,s) = L(H^2_{et}(\bar{X}, \mathbb{Q}_\ell), s) = L(NS(X) \otimes \mathbb{Q}_\ell, s) \times L(T(X) \otimes \mathbb{Q}_\ell, s).$$

We know that for a good prime p,

$$P_p^0(T) = 1 - T, P_p^4(T) = 1 - p^2 T,\ P_p^1(T) = P_p^3(T) = 1$$

and if

$$P_p^2(T) = \prod_{j=1}^{22}(1 - \alpha_j\, T) \in \bar{\mathbb{Q}}[T]$$

then

$$|\alpha_j| = p.$$

The L-series of $NS(X)$ is more or less understood by Tate's conjecture. The validity of the Tate conjecture for K3 surfaces in characteristic zero has been established (see Tate [59]). In fact, if we know that all the algebraic cycles generating $NS(X)$ are defined over some finite extension $\mathbb{K}$ of $\mathbb{Q}$ of degree r, then $\rho^2(\mathrm{Fr}_{p^r})$ acts on $NS(X) \otimes \mathbb{Q}_\ell$ by multiplication by p^r so that the L-series may be expressed as

$$L(NS(X_{\mathbb{K}}) \otimes \mathbb{Q}_\ell, s) = \zeta_{\mathbb{K}}(s-1)^{\rho(X)}$$

where $\zeta_{\mathbb{K}}(s)$ denotes the Dedekind zeta-function of $\mathbb{K}$.

Therefore, the remaining task is to determine the L-series $L(T(X) \otimes \mathbb{Q}_\ell, s)$ arising from the transcendental cycles $T(X)$, and we call this the motivic L-series of X.

(c) Let $d = 3$, and let X be a Calabi–Yau threefold defined over $\mathbb{Q}$. For a good prime p,

$$\#X(\mathbb{F}_p) = \sum_{i=0}^{6}(-1)^i t_i(p) = t_0(p) - t_1(p) + t_2(p) - t_3(p) + t_4(p) - t_5(p) + t_6(p)$$

$$= 1 + p^3 + (1+p)t_2(p) - t_3(p).$$

Therefore,

$$t_3(p) = 1 + p^3 + (1+p)t_2(p) - \#X(\mathbb{F}_p).$$

We have

$$t_0(p) = 1,\ t_6(p) = p^3,\ t_1(p) = t_5(p) = 0,$$

$$|t_2(p)| \le p h^{1,1}(X),\ t_4(p) = p t_2(p),$$

and

$$|t_3(p)| \le B_3\, p^{3/2}.$$

Hence

$$P_p(T) = 1 - T,\ P_p^6(T) = 1 - p^3 T,\ P_p^1(T) = P_p^5(T) = 1,$$

$$P_p^4(T) = P_2^2(pT) \quad \text{and} \quad P_p^3(T) \in \mathbb{Z}[T] \quad \text{with degree } B_3(X).$$

Then the L-series of X is given by

$$L(X, s) = L(H_{et}^3(\bar{X}, \mathbb{Q}_\ell), s) = (*) \prod_{p\ good} P_p^3(p^{-s})^{-1}$$

where $(*)$ is the factor corresponding to bad primes.

We will now make a definition of what it means for a Calabi–Yau variety X defined over $\mathbb{Q}$ to be modular (or automorphic). This is a concrete realization of the conjectures known as the Langlands Philosophy.

In the Appendix, we will briefly recall the definitions of various types of modular forms.

Now we will recall the Fontaine–Mazur conjectures, or rather some variant concentrated on Calabi–Yau varieties over $\mathbb{Q}$.

Definition 1.4 Let X be a Calabi–Yau variety of dimension $d \leq 3$ defined over $\mathbb{Q}$. Let $L(X,s)$ be its L-series. We say that X is modular if there is a set of modular forms (or automorphic forms) such that $L(X,s)$ coincides with the L-series associated to modular forms (automorphic forms), up to a finite number of Euler factors.

Remark 1.5 In fact, when the Galois representation arising from X is reducible, then we will consider its irreducible factors and match their L-series with the L-series of modular forms. This is the so-called motivic modularity.

Now we define Calabi–Yau varieties X of *CM-type*. This involves a polarized rational Hodge structure h on the primitive cohomology $H^d_{\text{prim}}(X,\mathbb{Q})$, where $d = \dim(X)$.

Let $\mathbb{S} := R_{\mathbb{C}/\mathbb{R}}\mathbb{G}_m$ be the real algebraic group obtained from $\mathbb{G}_m$ by restriction of scalars from $\mathbb{C}$ to $\mathbb{R}$. For $d=1$, $H^1_{\text{prim}}(X,\mathbb{Q}) = H^1(X,\mathbb{Q})$, and polarized rational Hodge structures on X are simple. For $d=2$, The Hodge group of a polarized rational Hodge structure $h : \mathbb{S} \to GL(H^d_{\text{prim}}(X,\mathbb{Q}) \otimes \mathbb{R})$ is the smallest algebraic group of $GL(H^2_{\text{prim}}(X,\mathbb{Q}) \otimes \mathbb{R})$ defined over $\mathbb{Q}$ such that the real points $\text{Hdg}(\mathbb{R})$ contain $h(U^1)$ where $U^1 := \{\, z \in \mathbb{C}^* \mid z\overline{z} = 1\}$. For details about Hodge groups, see Deligne [10], and Zarhin [66]. For $d = 3$, the Hodge structure on $H^3_{\text{prim}}(X,\mathbb{Q}) = H^3(X,\mathbb{Q})$ is not simple for all Calabi–Yau threefolds X. (For instance, for the quintic threefold in the Dwork pencil, the Hodge structure would split into 4-dimensional Hodge substructures, using the action of the $(\mathbb{Z}/5\mathbb{Z})^3$.)

Definition 1.6 A Calabi–Yau variety X of dimension $d \leq 3$ is said to be of CM *type* if the Hodge group $\text{Hdg}(X)$ associated to a rational Hodge structure of weight k of $H^d_{\text{prim}}(X,\mathbb{Q})$ is commutative. That is, the Hodge group $\text{Hdg}(X)_{\mathbb{C}}$ is isomorphic to a copy of $\mathbb{G}_m \simeq \mathbb{C}^*$.

Hodge groups are very hard to compute in practice. Here are algebraic characterizations of CM type Calabi–Yau varieties of dimension d.

Proposition 1.7 • $d = 1$. *An elliptic curve E over $\mathbb{Q}$ is of CM type if and only if* $\text{End}(E) \otimes \mathbb{Q}$ *is an imaginary quadratic field over $\mathbb{Q}$.*

• $d = 2$. *A K3 surface X over $\mathbb{Q}$ is of CM type if* $\text{End}_{\text{Hdg}}(T(X)) \otimes \mathbb{Q}$ *is a CM field over $\mathbb{Q}$ of degree equal to* $\operatorname{rank} T(X)$.

• $d = 3$. *Let X be a Calabi–Yau threefold over $\mathbb{Q}$. A Calabi–Yau threefold X over $\mathbb{Q}$ is of CM type if and only if* $\text{End}_{\text{Hdg}}(X) \otimes \mathbb{Q}$ *is a CM field over $\mathbb{Q}$ of degree* $2(1 + h^{2,1}(X))$*, if and only if the Weil and Griffiths intermediate Jacobians of X are of CM type.*

For $d = 1$, this is a classical result. For $d = 2$, a best reference might be Zarhin [66], and for $d = 3$, see Borcea [3].

Later we will construct Calabi–Yau varieties of dimension 2 and 3 which are of CM type.

2 Results on modularity of Galois representations

2.1 Two-dimensional Galois representations arising from Calabi–Yau varieties over $\mathbb{Q}$. We will focus on 2-dimensional Galois representations arising from Calabi–Yau varieties over $\mathbb{Q}$.

First, for dimension 1 Calabi–Yau varieties over $\mathbb{Q}$, we have the celebrated theorem of Wiles et al.

Theorem 2.1 *($d=1$) Every elliptic curve E defined over $\mathbb{Q}$ is modular. More concretely, let E be an elliptic curve over $\mathbb{Q}$ with conductor N. Then there exists a Hecke eigen newform f of weight $2=1+d$ on the congruence subgroup $\Gamma_0(N)$ such that*

$$L(E,s)=L(f,s).$$

That is, if we write $f(q)=\sum_{m=1}^{\infty}a_f(m)q^m$ with $q=e^{2\pi iz}$ and normalized by $a_f(1)=1$, then $a(n)=a_f(n)$ for every n.

For dimension 2 Calabi–Yau varieties, namely, K3 surfaces, over $\mathbb{Q}$, there are naturally associated Galois representations. In particular, for a special class of K3 surfaces, the associated Galois representations are 2-dimensional. Let X be a K3 surface defined over $\mathbb{Q}$ with Picard number $\rho(X_{\bar{\mathbb{Q}}})=20$. Such K3 surfaces are called *singular* K3 surfaces. Then the group (or lattice) $T(X)$ of transcendental cycles on X is of rank 2, and it gives rise to a 2-dimensional Galois representations. Livné [41] has established the motivic modularity of X, that is, the modularity of $T(X)$.

Theorem 2.2 *($d=2$) Let X be a singular K3 surface defined over $\mathbb{Q}$. Then $T(X)$ is modular, that is, there is a modular form f of weight $3=1+d$ on some $\Gamma_1(N)$ or $\Gamma_0(N)$ with a character ε such that*

$$L(T(X)\otimes\mathbb{Q}_\ell,s)=L(f,s).$$

Remark 2.3 A representation theoretic formulation of the above theorem is given as follows. Let π be the compatible family of 2-dimensional ℓ-adic Galois representations associated to $T(X)$ and let $L(\pi,s)$ be its L-series. Then there exists a unique, up to isomorphism, modular form of weight 3, level=conductor of π, and Dirichlet character $\varepsilon(p)=\left(\frac{-d}{p}\right)$ such that

$$L(\pi,s)=L(f,s).$$

Here $d=|\text{disc } NS(X)|$.

For dimension 3 Calabi–Yau varieties over $\mathbb{Q}$, we will focus on rigid Calabi–Yau threefolds. A Calabi–Yau threefold X is said to be *rigid* if $h^{2,1}(X)=0$ so that $B_3(X)=2$. This gives rise to a 2-dimensional Galois representation.

Theorem 2.4 *($d=3$) Every rigid Calabi–Yau threefold X over $\mathbb{Q}$ is modular. That is, there exists a cusp form f of weight $4=1+d$ on some $\Gamma_0(N)$ such that*

$$L(X,s)=L(f,s).$$

This theorem has been established by Gouvêa and Yui [26], and independently by Dieulefait [11]). Their proof relies heavily on the recent results on the modularity of Serre's conjectures about 2-dimensional residual Galois representations by Khare–Wintenberger [31] and Kisin [32].

A list of rigid Calabi–Yau threefolds over $\mathbb{Q}$ can be found in the monograph of Meyer [46].

2.2 Modularity of higher dimensional Galois representations arising from K3 surfaces over $\mathbb{Q}$. We will first consider K3 surfaces X with $T(X)$ of rank ≥ 3.

The first result is for K3 surfaces with transcendental rank 3.

Theorem 2.5 *Let X be a K3 surface over $\mathbb{Q}$ with Picard number 19. Then X has a Shioda–Inose structure, that is, X has an involution ι such that X/ι is birational to a Kummer surface Y over $\mathbb{C}$.*

Suppose that the Kummer surface is given by the product $E \times E$ of a non-CM elliptic curve E over $\mathbb{Q}$. Then the Shioda–Inose structure induces an isomorphism of integral Hodge structures on the transcendental lattices, so, X and $Km(E\times E)$ have the same $\mathbb{Q}$-Hodge structure. In this case, the 2-dimensional Galois representation ρ_E associated to E induces the 3-dimensional Galois representation $Sym^2\rho_E$ on $T(X)$ over some number field. Consequently, $T(X)$ is **potentially modular** *in the sense that the L-series of $T(X)$ is determined over some number field K by the symmetric square of a modular form g of weight 2 associated to E, and over K,*

$$L(T(X)\otimes \mathbb{Q}_\ell, s) = L(Sym^2(g), s).$$

Remark 2.6 In the above theorem, we are not able to obtain the modularity results over $\mathbb{Q}$. Since the K3 surface S is defined over $\mathbb{Q}$, the representation on $T(X)$ is also defined over $\mathbb{Q}$, but the isomorphism to $\mathrm{Sym}^2\rho_E$ may not be. Thus, we only have the potential modularity of X.

Can we say anything about the modularity of K3 surfaces with arbitrary large transcendental rank? The Galois representations associated to these K3 surfaces have large dimensions. The only result along this line is due to Livné-Schütt-Yui [42].

Consider a K3 surface X with non-symplectic automorphism. Let ω_X denote a holomorphic 2-form on X, fixed once and for all. Then $H^{2,0}(X) \simeq \mathbb{C}\omega_X$. Let $\sigma \in \mathrm{Aut}(X)$. Then σ induces a map

$$\sigma^* : H^{2,0}(X) \to H^{2,0}(X) \quad \omega_X \to \alpha\omega_X,\, \alpha \in \mathbb{C}^*.$$

We say that σ^* is non-symplectic if $\alpha \neq 1$. Let

$$H_X := \mathrm{Ker}(\mathrm{Aut}(X) \to O(NS(X)).$$

Then H_X is a finite cyclic group, and in fact, can be identified with the group of roots of unity μ_k for some $k \in \mathbb{N}$. Assume that $\det(T(X)) = \pm 1$ (that is, assume that $T(X)$ is unimodular.) Then we have the following possibilities for the values of k.

- $k \leq 66$ (Nikulin [49])
- k is a divisor of $66, 44, 42, 36, 28, 12$ (Kondo [33])

• If $\operatorname{rank}(T(X)) = \phi(k)$ (where ϕ is the Euler function), then $k = 66, 44, 42, 36, 28, 12$. Furthermore, there is a unique K3 surface X with given k. (Kondo [33]) These results were first announced by Vorontsov [62], and proofs were given later by Nikulin [49] and Kondo [33].

We tabulate these six K3 surfaces.

k	$NS(X)$	$T(X)$	$\operatorname{rank}(T(X))$
12	$U \oplus (-E_8)^2$	U^2	4
28	$U \oplus (-E_8)$	$U^2 \oplus (-E_8)$	12
36	$U \oplus (-E_8)$	$U^2 \oplus (-E_8)$	12
42	$U \oplus (-E_8)$	$U^2 \oplus (-E_8)$	12
44	U	$U^2 \oplus (-E_8)^2$	20
66	U	$U^2 \oplus (-E_8)^2$	20

Here $(-E_8)$ denotes the negative definite even unimodular lattice of rank 8.

Now we ought to realize these K3 surfaces over $\mathbb{Q}$. Here are explicit equations thanks to Kondo [33].

k	X	σ
12	$y^2 = x^3 + t^5(t^2+1)$	$(x, y, t) \mapsto (\zeta_{12}^2 x, \zeta_{12}^3 y, -t)$
28	$y^2 = x^3 + x + t^7$	$(x, y, t) \mapsto (-x, \zeta_{28}^7 y, \zeta_{28}^2 t)$
36	$y^2 = x^3 - t^5(t^6-1)$	$(x, y, t) \mapsto (\zeta_{36}^2 x, \zeta_3 6^3 y, \zeta_{36}^{30} t)$
42	$y^2 = x^3 + t^5(t^7-1)$	$(x, y, t) \mapsto (\zeta_{42}^2 x, \zeta_{42}^3 y, \zeta_{42}^{18} t)$
44	$y^2 = x^3 + x + t^{11}$	$(x, y, t) \mapsto (-x, \zeta_{44}^{11} y, \zeta_{44}^2 t)$
66	$y^2 = x^3 + t(t^{11}-1)$	$(x, y, t) \mapsto (\zeta_{66}^2 x, \zeta_{66}^3 y, \zeta_{66}^6 t)$

Here ζ_k denotes a primitive k-th root of unity.

Now the main result of Livné–Schütt–Yui [42] is to establish the modularity of these K3 surfaces.

Theorem 2.7 *Let X be a K3 surface in the above table. Then for each k, the ℓ-adic Galois representation associated to $T(X)$ is irreducible over $\mathbb{Q}$ of dimension $\phi(k)$. Furthermore, this $G_{\mathbb{Q}}$-Galois representation is induced from a one-dimensional Galois representation of $\mathbb{Q}(\zeta_k)$.*

All these K3 surfaces are of CM type, and are modular (automorphic).

Proof • CM type is established by realizing them as Fermat quotients. Since Fermat surfaces are known to be of CM type, the result follows.

• Modularity (or automorphy) of the Galois representation is established by using automorphic induction. The restriction of the Galois representation to the cyclotomic field $\mathbb{Q}(\mu_k)$ is given by a one-dimensional Jacobi sum Grössencharakter. To get down to $\mathbb{Q}$, we take the $\operatorname{Gal}(\mathbb{Q}(\zeta_k)/\mathbb{Q})$-orbit of the one-dimensional representation. This Galois group has order $\phi(k)$, and we obtain the irreducible Galois representation over $\mathbb{Q}$ of dimension $\phi(k)$.

• Those K3 surfaces corresponding to $k = 44$ and 66 are singular, so their modularity has already been established by Theorem 2.2.

Remark 2.8 When $T(X)$ is not unimodular, there are 10 values of k such that $\operatorname{rank} T(X) = \phi(k)$:

$$19, 17, 13, 11, 7, 25, 5, 27, 9, 3$$



Figure 1 Nikulin's pyramid

All these K3 surfaces are again dominated by Fermat surfaces, and hence they are all of CM type. We have also established their modularity (automorphy).

In the article of Goto–Livné–Yui [23], more examples of K3 surfaces of CM type are constructed. First we recall a classification result of Nikulin [50].

Let X be a K3 surface over $\mathbb{Q}$. Let $H^{2,0}(X) = \mathbb{C}\omega_X$ where we fix a nowhere vanishing holomorphic 2-form ω_X. Let σ be an involution on X such that $\sigma(\omega_X) = -\omega_X$. Let $= \operatorname{Pic}(X)^\sigma$ be the fixed part of $\operatorname{Pic}(X)$ by σ. Put $r = \operatorname{rank}\operatorname{Pic}(X)^\sigma$. Let $T(X)_0$ be the orthogonal complement of $\operatorname{Pic}(X)^\sigma$ in $H^2(X,\mathbb{Z})$. Then σ acts by -1 on $T(X)_0$. Consider the quotient groups $(\operatorname{Pic}(X)^\sigma)^*/\operatorname{Pic}(X)^\sigma)$ and $(T(X)_0^*/T(X)_0)$, where L^* denotes the dual lattice of a lattice L. Since $H^2(X,\mathbb{Z})$ is unimodular, the quotient abelian groups are canonically isomorphic

$$(\operatorname{Pic}(X)^\sigma)^*/\operatorname{Pic}(X)^\sigma \simeq (T(X)_0^*/T(X)_0.$$

Since σ acts as 1 on the first quotient; and as -1 on the second quotient, this forces these quotient groups to be isomorphic to $(\mathbb{Z}/2\mathbb{Z})^a$ for some positive integer $a \in \mathbb{Z}$.

The intersection pairing on $\operatorname{Pic}(X)$ induces a quadratic form q on the discriminant group with values in $\mathbb{Q}$ modulo $2\mathbb{Z}$; we put $\delta = 0$ if q has values only in $\mathbb{Z}$, and $\delta = 1$ otherwise.

Thus, we have a triplet of integers (r, a, δ) associated

to a K3 surface X with the involution σ. A theorem of Nikulin [50] asserts that a pair (X, σ) is classified, up to deformation, by a triplet (r, a, δ).

Theorem 2.9 *There are* 75 *possible triplets* (r, a, δ) *that classify pairs* (X, σ) *of K3 surfaces* X *with involution* σ*, up to deformation.*

Now we will realize some of these 75 families of K3 surfaces with involution. We look for K3 surfaces defined by hypersurfaces. For this we use the famous 95 families of hypersurfaces in weighted projective 3-spaces determined by M. Reid [52] or Yonemura [63]. Let $[x_0 : x_1 : x_2 : x_3]$ denote weighted projective coordinates in a weighted projective 3-space with weight (w_0, w_1, w_2, w_3).

Theorem 2.10 *(Goto–Livné–Yui* [23]*) Among the* 95 *families of K3 surfaces, all but* 9 *families of K3 surfaces have an involution* σ*, satisfying the following conditions:*

(1) removing several monomials (if necessary) from Yonemura's hypersurface, a new defining hypersurface consists of exactly four monomials, i.e., it is of Delsarte type,

(2) the new defining hypersurface is quasi-smooth, and the singularity configuration should remain the same as the original defining equation of Yonemura,

(3) the new defining hypersurface contains only one monomial in x_0 *of the form* x_0^n, $x_0^n x_j$, $x_i^n + x_i x_j^m$ *or* $x_i^n x_k + x_i x_j^m$ *for some* j *and* $k\,(k \neq j)$ *distinct from* i*.*

For 45 *(resp.* 41*) K3 surfaces, a defining equation of four monomials is of the form*

$$x_0^2 = f(x_1, x_2, x_3) \subset \mathbb{P}^3(w_0, w_1, w_2, w_3)$$

(resp.

$$F(x_0, x_1, x_2, x_3) = 0 \subset \mathbb{P}^3(w_0, w_1, w_2, w_3))$$

where f *(resp.* F*) is a homogeneous polynomial in* x_1, x_2, x_3 *(resp.* x_0, x_1, x_2, x_3*) over* $\mathbb{Q}$ *of degree* $\sum_{i=0}^{3} w_i$*.*

(4) For all $86 = 45 + 41$ *K3 surfaces, there is an algorithm to compute the invariants* r *and* a*.*

Remark 2.11 There are nine families for which the Theorem is not valid. The three families (#15, #53 and #54) do not have the required involution. Another different six families (#85, #94, #95) and (#90, #93, #91) cannot not realized as quasi-smooth hypersurfaces in four monomials. (We employ the numbering from Yonemura [63].)

Proposition 2.12 *(Nikulin* [50]*; Voisin* [61]*) Let* (S, σ) *be a K3 surface with involution* σ*. Let* S^σ *be the fixed part of* S*. Then for* $(r, a, \delta) \neq (10, 10, 0), (10, 8, 0)$*, let* $S^\sigma = C_g \cup L_1 \cup \cdots \cup L_k$ *where* C_g *is a smooth genus* g *curve and* $L_1, \ldots, L_k$ *are rational curves. Then*

$$r = 11 + k - g, \; a = 11 - g - k.$$

If $(r, a, \delta) = (10, 10, 0)$*, then* $S^\sigma = C_1 \cup C_2$ *where* C_i $(i = 1, 2)$ *are elliptic curves, and if* $(r, a, \delta) = (10, 8, 0)$*, then* $S^\sigma = \emptyset$*.*

Proposition 2.13 *(Goto–Livné–Yui* [23]*) Let* (S, σ) *be one of the* 45 *pairs in the above theorem, which is given as the minimal resolution of a hypersurface* $S_0 : x_0^2 = f(x_1, x_2, x_3) \subset \mathbb{P}^3(Q)$ *where* $Q = (w_0, w_1, w_2, w_3)$ *and* f *is a homogeneous polynomial of degree* $\sum_{i=0}^{3} w_i$*. Let* $r(Q)$ *denote the number of exceptional divisors in the resolution* $S \to S_0$*. Then* r *can be computed as follows:*

(i) Suppose w_0 *is odd. Then* $r = r(Q) - w_i + 2$ *if there is an odd weight* $w_i \neq w_0$ *such that* $\gcd(w_0, w_i) = w_i \geq 2$*, and* $r = r(Q) + 1$ *otherwise.*

(ii) Suppose w_0 *is even. Then* $r = r(Q) + 1 - \sum_{i=1}^{3}(d_i - 1)(\frac{2d_i}{w_i} - 1)$ *where* $d_i = \gcd(w_0, w_i)$*.*

Remark 2.14 For non-Borcea type K3 surfaces defined by equations of the form $x_0^2 x_i = f(x_1, x_2, x_3)$ for some $i \in \{1, 2, 3\}$, the invariants r and a (or equivalently, g and k) are also computed.

Proposition 2.15 *(Goto–Livné–Yui* [23]*) At least the* 39 *triplets* (r, a, δ) *of integers are realized by the* 86 *families of K3 surfaces.*

Theorem 2.16 *(Goto–Livné–Yui* [23]*) For the* 86 *families of K3 surfaces, there are subfamilies of K3 surfaces which are of CM type, that is, there is a CM point in the moduli space of K3 surfaces. Indeed, at each CM point, the K3 surface is realized as a quotient of a Fermat (or Delsarte) surface. Consequently, it is modular (automorphic) by automorphic induction.*

Remark 2.17 Nikulin's Theorem 2.9 gives in total 75 triplets of integers (r, a, δ) that classify the isomorphism classes of pairs (S, σ) of K3 surfaces S with non-symplectic involution σ. With our choice of K3 surfaces of CM type, we can realize at least 39 triplets. This is based on our calculations of the invariants r and a. Since we have not yet computed the invariant δ for the 39 cases, this number may increase somewhat.

Remark 2.18 It is known (Borcea [4]) that over $\mathbb{C}$ the moduli spaces of Nikulin's K3 surfaces are arithmetic quotients of type IV (Shimura varieties). Recently, Ma [45] has shown the rationality of the moduli spaces for the 67 triplets. Our result above gives one explicit CM point in these moduli spaces. CM type implies that these families must be isolated points in the moduli space. We are not able to show the denseness of CM points, however.

It is notoriously difficult to compute Hodge groups, and the above theorem implies the commutativity of the Hodge group.

Corollary 2.19 The Hodge groups of these 86 K3 surfaces are all commutative, i.e., copies of $\mathbb{G}_m$'s over $\mathbb{C}$.

Proposition 2.20 *(Goto-Livné–Yui* [23]*) We classify the* 86 *hypersurfaces into four types:*

- 45 *families have the form:* $x_0^2 = f(x_1, x_2, x_3)$ *and* $\sigma(x_0) = -x_0$.
- 33 *families cannot be put in the form (1) but defined by a hypersurface of the form* $F(x_0, x_1, x_2, x_3) = 0$ *with* 4 *monomials, and* σ *can be described explicitly.*
- 4 *families, after changing term the* x_0^3 *to* $x_0^2x_1$ *and then removing several terms, can be put into the form* $F(x_0, x_1, x_2, x_3) = 0$ *of four monimials equipped with an explicit involution* σ.
- *The remaining* 4 *families can be put in the form* $F(x_0, x_1, x_2, x_3) = 0$ *of four monomials equipped with a different kind of involution.*

Example 2.21 (1) Here is one of the 45 cases, #78 in Yonemura's list. The weight is $(11, 6, 4, 1)$, and Yonemura's hypersurface is

$$x_0^2 = x_1^2x_2 + x_1^3x_3^4 + x_1x_2^4 + x_2^5x_3^2 + x_3^{22}.$$

We can remove $x_1^3x_2^4$ and $x_2^5x_3^2$ so the new hypersurface is $x_0^2 = x_1^2x_2 + x_1x_2^4 + x_3^{22}$. The singularity is of type $A_1 + A_3 + A_5$.

(2) Here is one of the second cases, #19 in Yonemura's list. The weight is $(3, 2, 2, 1)$ and Yonemura's hypersurface is

$$F(x_0, x_1, x_2, x_3) = x_0^2x_1 + x_0^2x_2 + x_0^2x_3^2 + x_1^4 + x_2^4 + x_3^4.$$

We can remove $x_0^2x_1$ or $x_0^2x_2$, $x_0^2x_3^2$, so the new hypersurface is $x_0^2x_1 + x_1^4 + x_2^4 + x_3^4$. The singularity is of type $4A_1 + A_2$.

(3) Here is one of the third cases, #18 in Yonemura's list. This hypersurface $x_0^3 + x_1^3 + x_0x_2^3 + x_1x_2^3 + x_2^4x_3 + x_3^9$ acquires an involution if we replace x_0^3 by $x_0^2x_1$ and remove the terms $x_0x_2^3$ and $x_1x_2^3$.

(4) The last case is #52 in Yonemura's list.

(5) #95 in Yonemura's list. This hypersurface has an involution, but cannot be realized by a quasi-smooth hypersurface with four monomials.

2.3 The modularity of higher dimensional Galois representations arising from Calabi–Yau threefolds over $\mathbb{Q}$. There are several new examples of modular non-rigid Calabi–Yau threefolds X over $\mathbb{Q}$ with $B_3 \geq 4$. These examples were constructed after the article Yui[65], some of which have already been discussed in the article of E. Lee [37].

There are several approaches (with non-empty intersection) to produce these new modular examples:

(1) Those non-rigid Calabi–Yau threefolds X over $\mathbb{Q}$ such that the semi-simplification of $H^3_{et}(\bar{X}, \mathbb{Q}_\ell)$ is highly reducible and splits into smaller dimensional irreducible Galois representations. For instance, most known cases are when the third cohomology group splits into 2-dimensional or 4-dimensional pieces. (Examples of E. Lee, Hulek and Verrill, Schütt, Cynk and Meyer, and a recent example of Bini and van Geemen, and Schütt, and more.) The article of E. Lee [37] gives reviews on the modularity of non-rigid Calabi–Yau threefolds over $\mathbb{Q}$, up to 2008.

(2) Those Calabi–Yau threefolds X over $\mathbb{Q}$ such that the the ℓ-adic Galois representations arising from $H^3(\bar{X}, \mathbb{Q}_\ell)$ are irreducible and have small dimensions (e.g., $4, 6$, or 8.) (Examples of Livné–Yui, Dieulefeit–Pacetti–Schütt, and more.)

(3) Those Calabi–Yau threefolds over $\mathbb{Q}$ which are of CM type, thus the Galois representations are induced by one-dimensional representations. (Examples of Rohde, Garbagnati–van Geemen, Goto–Livné–Yui, and more.)

(4) Given a Calabi–Yau threefold Y over $\mathbb{Q}$, if we can construct an algebraic correspondence defined over $\mathbb{Q}$ to some modular Calabi–Yau threefold X over $\mathbb{Q}$, the Tate conjecture asserts that their L-series should coincide. This will establish the modularity of Y.

(1): Calabi–Yau threefolds in the category (1), i.e, with highly reducible Galois representations

The general strategy is to consider Calabi–Yau threefolds which contain elliptic ruled surfaces. This is formulated by Hulek and Verrill [29].

Proposition 2.22 *(Hulek and Verrill* [29]*) Let X be a Calabi–Yau threefold over $\mathbb{Q}$. Suppose that X contains birational ruled elliptic surfaces S_j, $j = 1, \ldots, b$ over $\mathbb{Q}$ and whose cohomology classes span $H^{2,1}(X) \oplus H^{1,2}(X)$ (so $b = h^{2,1}(X)$.) Let ρ be the 2-dimensional Galois representation given by the kernel U from the exact sequence*

$$0 \to U \to H^3_{et}(\bar{X}, \mathbb{Q}_\ell) \to \oplus H^3_{et}(\bar{S}_j, \mathbb{Q}_\ell) \to 0.$$

Then X is modular, that is,

$$L(X, s) = L(f_4, s) \prod_{j=1}^{b} L(g_2^j, s-1)$$

where f_4 is a weight 4 modular form associated to ρ and g_2^j are the weight 2 modular forms associated to the base elliptic curves E_j of the birational ruled surfaces S_j.

The requirement that the third cohomology group splits as in the proposition is rather restrictive. Several examples of Calabi–Yau threefolds satisfying this condition are given by Hulek and Verrill [29], E. Lee [36], and Schütt [56]. They are constructed as resolutions of fiber products of semi-stable rational elliptic surfaces with section. Another series of examples along this line due to Hulek and Verrill [30] are toric Calabi–Yau threefolds associated to the root lattice A_4.

Cynk and Meyer [9] have established the modularity of 17 nonrigid double octic Calabi–Yau threefolds over $\mathbb{Q}$ with $B_3 = 6$. The Galois representations decompose into 2- and 4-dimensional sub-representations such that the L-series of each such sub-representation is of the form $L(g_4, s)$, $L(g_2, s-1)$ or $L(g_2 \times g_3, s)$, where g_k is a weight k cusp form.

Now we will consider another construction due to E. Lee [36, 38] of Calabi–Yau threefolds associated to the Horrocks–Mumford vector bundle of rank 2. It is well known that Horrocks–Mumford quintics are determinantal quintics. The Schoen quintic $Q : \sum_{i=0}^4 x_i^5 - 5\prod_{i=0}^4 x_i = 0$ is the early example of this type.

Lee has constructed more Calabi–Yau threefolds. Let $y \in \mathbb{P}^4$ be a generic point, and define the matrices

$$M_y(x) = \begin{pmatrix} x_0y_0 & x_3y_2 & x_1y_4 & x_4y_1 & x_2y_3 \\ x_3y_3 & x_1y_0 & x_4y_2 & x_2y_4 & x_0y_1 \\ x_1y_1 & x_4y_3 & x_2y_0 & x_0y_2 & x_3y_4 \\ x_4y_4 & x_2y_1 & x_0y_3 & x_3y_0 & x_1y_2 \\ x_2y_2 & x_0y_4 & x_3y_1 & x_1y_3 & x_4y_0 \end{pmatrix}$$

and

$$L_y(z) = \begin{pmatrix} z_0y_0 & z_2y_4 & z_4y_3 & z_1y_2 & z_3y_1 \\ z_4y_1 & z_1y_0 & z_3y_4 & z_0y_3 & z_2y_2 \\ z_3y_2 & z_0y_1 & z_2y_0 & z_4y_4 & z_1y_3 \\ z_2y_3 & z_4y_2 & z_1y_1 & z_3y_0 & z_0y_4 \\ z_1y_4 & z_3y_3 & z_0y_2 & z_2y_1 & z_4y_0 \end{pmatrix}$$

Note that $M_y(x)z = L_y(z)x$. Then

$$X_y := \{\det M_y(x) = 0\} \subset \mathbb{P}^4(x)$$

and

$$X'_y := \{\det L_y(z) = 0\} \subset \mathbb{P}^4(z)$$

are Horrocks–Mumford quintics. Define a threefold $\tilde{X}_y$ in $\mathbb{P}^4(x) \times \mathbb{P}^4(z)$ as a common partial resolution of $M_y(x)z = 0$. We ought to know the singularities of $\tilde{X}_y$. Let $\mathbb{P}^2_+ := \{y : y_1 - y_4 = y_2 - y_3 = 0\}$. Given $y \in \mathbb{P}^2_+$, the point $(1:0:0:0:0)$ is a singular point over $\mathbb{C}$ of X_y if and only if one of the coordinates of y is zero. For $y = (0:1:-1:-1:1:1), (0:2:3 \pm \sqrt{5}:3 \pm \sqrt{5}:2), (2:-1:0:0:-1), (2:\pm\sqrt{5}-1:0:0:\pm\sqrt{5}-1) \in \mathbb{P}^2_+$, X_y contains the Heisenberg orbits of $(1:0:0:0:0)$ and $(1:1:1:1:1)$ as nodes over $\mathbb{C}$.

Proposition 2.23 *(Lee* [36, 37, 38]*) (a) Let* $y = (0:1:-1:-1:1)$*, and write* $X_{(0:1:-1:-1:1)} = X$ *for short. Then the Calabi–Yau threefold* $\hat{X}$ *obtained by crepant resolution of singularities has* $B_3 = 6$*. Furthermore,*

$$H^3_{et}(\bar{\hat{X}}, \mathbb{Q}_\ell) = V \oplus H^2(\bar{S}, \mathbb{Q}_\ell)(-1)$$

where V *is* 2*-dimensional and associated to the modular form of weight* 4 *and level* 55*, and* $H^2(\bar{S},\mathbb{Q}_\ell)$ *is* 4*-dimensional and is isomorphic to* $Ind_{G_{\mathbb{Q}(i)}}^{G_{\mathbb{Q}}} H^1(\bar{E},\mathbb{Q}_\ell)(-1)$ *where* E *is an elliptic curve over* $\mathbb{Q}(i)$ *coming from* E *over* $\mathbb{Q}$*.*

(b) Let $y = (2 : -1 : 0 : 0 : -1)$*, and let* $X_{(-2:-1:0:0:-1)} = X'$ *for short. Then the Calabi–Yau threefold* $\hat{X}'$ *obtained by crepant resolution of singularities has* $B_3 = 4$*. Furthermore*

$$H^3_{et}(\bar{\hat{X}},\mathbb{Q}_\ell) = V \oplus H^1(\bar{E}_2,\mathbb{Q}_\ell)(-1)$$

where V *is the* 2*-dimensional Galois representation associated to the modular form of weight* 4 *of level* 55*, and* $H^1(\bar{E}_2,\mathbb{Q}_\ell)$ *is associated with the modular form of weight* 2 *and level* 550*. Furthermore, the L-series of* $\hat{X}$ *is given, up to Euler factors at the primes of bad reduction, by*

$$L(\hat{X},s) = L(f,s)L(g,s-1)$$

where f *is the unique normalized cusp form of weight* 4 *and level* 55 *and* g *is the normalized cusp form of weight* 2 *and level* 550*.*

(c) Now consider a smooth (big) resolution X *of the* $(\mathbb{Z}/2\mathbb{Z})$*-quotient of the Schoen quintic* $Q : x_0^5+x_1^5+x_2^5+x_3^5+x_4^5-5x_0x_1x_2x_3x_4 = 0$*. (There is the* $(\mathbb{Z}/2\mathbb{Z})$*-action on* Q *induced by the involution on* $\mathbb{P}^4$ *defined by* $\iota[x_0 : x_1 : x_2 : x_3 : x_4] = [x_0 : x_4 : x_3 : x_2 : x_1]$*.) Then* X *is a Calabi–Yau threefold defined over* $\mathbb{Q}$ *with* $B_3 = 4$*. The Calabi–Yau threefold* X *is modular, and up to Euler factors at primes of bad reduction at* $p = 2$ *and* 5*,* $L(X,s)$ *is given by*

$$L(X,s) = L(f,s)L(g,s-1)$$

where f *is the unique normalized cusp form of weight* 4 *and level* 25*, and* g *is a weight* 2 *cusp form of level* 50*.*

A recent example due to Bini and van Geemen [2], and Schütt [57] is the Calabi–Yau threefold called Maschke's double octic, which arises as the double covering of $\mathbb{P}^3$ branched along Maschke's surface S. The Maschke octic surface S is defined by the homogeneous equation

$$S = \sum_{i=0}^{3} x_i^8 + 14\sum_{i<j} x_i^4x_j^4 + 168x_0^2x_1^2x_2^2x_3^2 = 0 \subset \mathbb{P}^3$$

Now let X be the double cover of $\mathbb{P}^3$ along S. This is a smooth Calabi–Yau threefold defined over $\mathbb{Q}$. Let Y be the desingularization of the quotient of X by a suitable Hisenberg group. Then Y is also a smooth Calabi–Yau threefold defined over $\mathbb{Q}$. The results of Bini and van Geemen, and Schütt, are summarized in the following

Proposition 2.24 *(a) The Maschke double octic Calabi–Yau threefolds* X *is modular over* $\mathbb{Q}$*. The third cohomology group* $H^3_{et}(\bar{X},\mathbb{Q}_\ell)$ *has* $B_3(X) = 300$*. The Galois representation of* $H^3_{et}(\bar{X},\mathbb{Q}_\ell)$ *decomposes completely over* $\mathbb{Q}(i)$ *into* 2*–dimensional Galois representations which descend to* $\mathbb{Q}$*, and the latter correspond to modular forms of weight* 4*, or modular forms of weight* 2*.*

(b) Let Y *be the desingularization of* X *by a suitable Heisenberg group. Then* Y *is modular over* $\mathbb{Q}$*. The third cohomology group* $H^3_{et}(\bar{Y},\mathbb{Q}_\ell)$ *has* $B_3(Y) = 30$*. The Galois representation of* $H^3_{et}(\bar{Y},\mathbb{Q}_\ell)$ *decomposes completely over* $\mathbb{Q}$ *into* 2*-dimensional Galois representations, and the latter correspond to modular forms of weight* 4 *and modular forms of weight* 2*.*

(c) The Maschke surface S has Picard number $\rho(S) = 202$. The second cohomology group $H^2(\bar{S}, \mathbb{Q}_\ell)$ has $B_2(S) = 302$. The Galois representation of the transcendental part has dimension 100*, which splits into* 2 *or* 3*-dimensional Galois sub-representations over $\mathbb{Q}$, and the latter correspond to modular forms of weight* 3*, or modular forms of weight* 2*.*

(2) Calabi–Yau threefolds in the category (2), i.e, with irreducible Galois representations

Consani–Scholten [7] constructed a Calabi–Yau threefold over $\mathbb{Q}$ as follows. Consider the Chebyshev polynomial

$$P(y,z) = (y^5+z^5) - 5yz(y^2+z^2) + 5yz(y+z) + 5(y^2+z^2) - 5(y+z)$$

and define an affine variety X in $\mathbf{A}^4$ by

$$X : P(x_1, x_2) = P(x_3, x_4)$$

and let $\overline{X} \subset \mathbb{P}^4$ be its projective closure. Then $\overline{X}$ has 120 ordinary double points. Let $\tilde{X}$ be its small resolution. Then $\tilde{X}$ is a Calabi–Yau threefold with $h^{1,1}(\tilde{X}) = 141$ and $h^{2,1}(\tilde{X}) = 1$. $\tilde{X}$ is defined over $\mathbb{Q}$ and the primes $2, 3, 5$ are bad primes. $H^3(\bar{\tilde{X}}, \mathbb{Q}_\ell)$ gives rise to a 4-dimensional ℓ-adic Galois representation, ρ, which is irreducible over $\mathbb{Q}$. Let $F = \mathbb{Q}(\sqrt{5})$, and let $\lambda \in F$ be a prime above ℓ. Then the restriction $\rho|_{\mathrm{Gal}(\bar{\mathbb{Q}}/F)}$ is reducible as a representation to $GL(4, F_\lambda)$: There is a Galois representation $\sigma : \mathrm{Gal}(\bar{\mathbb{Q}}/F) \to GL(2, F_\lambda)$ such that $\rho = \mathrm{Ind}_F^{\mathbb{Q}}\sigma$. Consani–Scholten conjectured the modularity of $\tilde{X}$.

Theorem 2.25 *(Dieulefait–Pacetti–Schütt* [12]*) The Consani-Scholten Calabi–Yau threefold $\tilde{X}$ over $\mathbb{Q}$ is Hilbert modular. That is, the L-series associated to σ coincides with the L-series of a Hilbert modular newform $\mathfrak{f}$ on F of weight* $(2,4)$ *and conductor $\mathfrak{c}_\mathfrak{f} = (30)$.*

The first example of Siegel modular varieties, as moduli spaces, of Calabi–Yau threefolds was given by van Geemen and Nygaard [24]. Recently, a series of articles by Freitag and Salvati Manni [19] on Siegel modular threefolds which admit Calabi–Yau models have appeared. The starting point is the van Geemen–Nygaard rigid Calabi–Yau threefold defined by a complete intersection Y of degree $(2,2,2,2)$ in $\mathbb{P}^7$ by the equations:

$$\begin{array}{ccccccccc} Y_0^2 & = & X_0^2 & + & X_1^2 & + & X_2^2 & + & X_3^2 \\ Y_1^2 & = & X_0^2 & - & X_1^2 & + & X_2^2 & - & X_3^2 \\ Y_2^2 & = & X_0^2 & + & X_1^2 & - & X_2^2 & - & X_3^2 \\ Y_3^2 & = & X_0^2 & - & X_1^2 & - & X_2^2 & + & X_3^2 \end{array}$$

A smooth small resolution of Y, denoted by X, is a Calabi–Yau threefold with $h^{1,1}(X) = 32$ and $h^{2,1}(X) = 0$, so X is rigid and hence is modular, indeed, the L-series of the Galois representation associated to $H^3(X, \mathbb{Q}_\ell)$ is determined by the unique weight 4 modular form on $\Gamma_0(8)$.

Several examples of non-rigid Calabi–Yau threefolds Z are obtained by quotienting X by finite group actions. Many of the resulting varieties Z are Calabi–Yau threefolds with small third Betti number.

Remark 2.26 For instance, Freitag–Salvati Manni has constructed such Calabi–Yau threefolds Z. The Galois representations associated to $H^3(Z, \mathbb{Q}_\ell)$ of their Calabi–Yau threefolds Z should be studied in detail. They should decompose into direct sum of those coming from the rigid Calabi–Yau threefold X, and those coming from elliptic surfaces or surfaces of higher genus arising from fixed points of finite groups in question.

In particular, this would imply that proper Siegel modular forms will not arise from these examples. So far as I know, we do not have examples of Calabi–Yau threefolds over $\mathbb{Q}$ with $B_3(X) = 4$ whose $L(X, s)$ comes from a Siegel modular form on $Sp_4(\mathbb{Z})$ or its subgroups of finite index.

(3) Calabi–Yau threefolds of CM type

We next consider Calabi–Yau threefolds which we will show to be of CM type. Then the Galois sub-representations associated to the third cohomology groups are induced by one-dimensional ones. Then, by applying the automorphic induction process, we will establish the automorphy of Calabi–Yau threefolds. These Calabi–Yau threefolds are realized as quotients of products of K3 surfaces and elliptic curves by some automorphisms. Rohde [53], Garbagnati–van Geemen [21], and Goto–Livné–Yui [23] produce examples of CM type Calabi–Yau threefolds with this approach.

Let S be a K3 surface with an involution σ acting on $H^{0,2}(S)$ by -1 discussed in subsection 2.2. Let E be an elliptic curve with the standard involution ι. Consider the quotient of the product $E \times S/\iota \times \sigma$. This is a singular Calabi–Yau threefold having only cyclic quotient singularities. Resolving singularities we obtain a smooth crepant resolution X. Since the invariants of X are determined by a triplet of integers (r, a, δ) associated to S, we will write X as $X(r, a, \delta)$. The Hodge numbers and the Euler characteristic of X depend only on r and a:

$$h^{1,1}(X) = 5 + 3r - 2a,\ h^{2,1}(X) = 65 - 3r - 2a$$

and

$$e(X) = 2(h^{1,1}(X) - h^{2,1}(X)) = 6(r - 10).$$

Theorem 2.27 *(Goto–Livné–Yui* [23]*) Let* (S, σ) *be one of the K3 surfaces defined over* $\mathbb{Q}$ *in Theorem 2.9. Let* E *be an elliptic curve over* $\mathbb{Q}$ *with the standard involution* ι*. Let* $X = X(r, a, \delta)$ *be a smooth Calabi–Yau threefold. Then the following assertions hold:*

(1) X *is of CM type if and only if* E *is of CM type.*

(2) If X *is of CM type, then the Jacobian variety* $J(C_g)$ *of* C_g *in* S^σ *is also of CM type, provided that the K3 surface component is of the form* $x_0^2 = f(x_1, x_2, x_3)$ *with involution* $\sigma(x_0) = -x_0$*.*

(3) X *is modular (automorphic).*

Sketch of Proof: (1) The Hodge structure h_X of type $(3, 0)$ of X is given by the tensor product $h_S \otimes h_E$ of the Hodge structures h_S of type $(2, 0)$ and h_E of type $(1, 0)$. Then h_X is of CM type if and only if both h_S and h_E are of CM type. Since S is already of CM type, we only need to require that E is of CM type.

(2) When the K3 surface is defined by a hypersurface of the form $x_0^2 = f(x_1, x_2, x_3)$ and the involution σ takes x_0 to $-x_0$, then the curve C_g in the fixed locus S^σ is obtained by putting $x_0 = 0$, and hence it is also of Delsarte type. Hence the Jacobian variety $J(C_g)$ of C_g is also of CM type.

When the hypersurface defining the K3 surface is not of the above form and the involution is more complicated, we ought to check each case whether C_g *is of Delsarte type or not.*

(3) S is modular by Theorem 2.11, and E is modular by Wiles et al. Hence X is modular (automorphic).

Now we will discuss mirror Calabi–Yau threefolds of $X = X(r, a, \delta)$.

Proposition 2.28 *(Borcea* [4] *and Voisin* [61]*) Given a Calabi–Yau threefold* $X = X(r, a, \delta)$*, there is a mirror Calabi–Yau threefold* $X^\vee$ *such that* $X^\vee$ *is realized as a crepant resolution of a quotient of* $E \times S/\iota \times \sigma$ *and* $X^\vee$ *is characterized by the invariants* $(20 - r, a, \delta)$*. The Hodge numbers of* $X^\vee$ *are*

$$h^{1,1}(X^\vee) = 5 + 3(20 - r) - 2a = 65 - 3r - 2a = h^{2,1}(X),$$

$$h^{2,1}(X^\vee) = 65 - 3(20 - r) - 2a = 5 + 3r - 2a = h^{1,1}(X)$$

and the Euler characteristic is

$$e(X^\vee) = -12(r - 10) = -e(X).$$

In terms of g *and* k*,* $r = 11 - g + k$*,* $a = 11 - g - k$*, and*

$$h^{1,1}(X) = 1 + r + 4(k + 1),\ h^{2,1}(X) = 1 + (20 - r) + 4g$$

and the Euler characteristic is $e(X) = 12(1 + k - g)$*.*

Remark 2.29 Mirror symmetry of Calabi–Yau threefolds of K3 × E do come from mirror symmetry of K3 surfaces. Mirror symmetry for K3 surfaces is the correspondence $r \leftrightarrow (20 - r)$. In fact, one can see this in Figure 1: Nikulin's pyramid. Given a K3 surface S, we try to look for a mirror K3 surface $S^\vee$ satisfying this correspondence. This correspondence is established at a special CM point in the moduli space.

We know that the 95 families of K3 surfaces of Reid and Yonemura are not closed under mirror symmetry. Only 54 families of K3 surfaces with involution have mirror partners within the 95 families.

A recent article of Artebani–Boissière–Sarti [1] also considers this type of mirror symmetry for K3 surfaces.

Proposition 2.30 *For a mirror* $X^\vee$*, the assertion is true if the mirror K3 surface is also of Delsarte type.*

Proposition 2.31 *Let* (S, σ) *be one of the* 86 *families of K3 surfaces with involution* σ*. Let* $X = X(r, a, \delta)$ *and* $X^\vee = X(20-r, a, \delta)$ *be mirror pairs of Calabi–Yau threefolds defined over* $\mathbb{Q}$*, where we suppose that the K3 surface component is of Delsarte type. Then* X *and* $X^\vee$ *have the same properties:*

- X *is of CM type if and only if* $X^\vee$ *is of CM type, and*
- X *is modular if and only if* $X^\vee$ *is modular.*

Rohde [53], Garbagnati and van Geemen [21] and Garbagnati [20] constructed Calabi–Yau threefolds which are quotients of the products of K3 surfaces and elliptic curves by non-symplectic automorphisms of higher order (than 2), that is, order 3, or order 4. These Calabi–Yau threefolds are parametrized by Shimura varieties.

Rohde [53] constructed families of Calabi–Yau threefolds as the desingularization of the quotient $S \times E$ by an automorphism of order 3 where E is the unique elliptic curve with an automorphism α_E of order 3, and S is a K3 surface with an automorphism α_S of order 3 which fixes k rational curves and $k + 3$ isolated points for some integer k, $0 \leq k \leq 6$.

Let ξ be a primitive cube root of unity. Choose the specific elliptic curve: $E = \mathbb{C}/\mathbb{Z}+\xi\mathbb{Z}$, which has a Weierstrass model $y^2 = x^3 - 1$ and $\alpha_E : (x, y) \mapsto (\xi x, y)$. Also choose some specific K3 surface S. Let

$$S = S_f : Y^2 = X^3 + f(t)^2,\ f = gh^2,\ \deg(f) = 6$$

where t is the coordinate on $\mathbb{P}^1$, and f has four distinct zeros. S_f has an automorphism of order 3: $\alpha_f : (X, Y, t) \mapsto (\xi X, Y, t)$. Now define a Calabi–Yau threefold X_f as the desingularization of $S_f \times E$ by the automorphism $\alpha := \alpha_f \times \alpha_E$. Note that S_f is birationally isomorphic to $(C_f \times E)/(\beta_f \times \alpha_E)$ where $C_f : v^3 = f(t)$ and $\beta_f : C_f \to C_f,\ (t, v) \mapsto (t, \xi v)$. Then X_f is birationally isomorphic to

$$(C_f \times E \times E)/H \quad \text{where} \quad H =< \beta_f \times \alpha_E \times 1_E, 1_{C_f} \times \alpha_E \times \alpha_E^{-1} > .$$

Rohde [53] worked out the case $\deg(g) = \deg(h) = 2$. Garbagnati–van Geemen [21] considered the other cases, i.e., $\deg(g) = 4, \deg(h) = 1$ and $\deg(g) = 6, \deg(h) = 0$. The Hodge numbers of the Calabi–Yau threefold X_f are computed and the results are tabulated as follows:

$\deg(g)$	$\deg(h)$	$g(C_f)$	$h^{2,1}(X_f)$	$h^{1,1}(X_f)$	k
6	0	4	3	51	3
4	1	3	2	62	4
2	2	2	1	73	5
0	3	1	0	84	6

For $k > 2$, the Calabi–Yau threefold is constructed by considering the curve $C_\ell : v^6 = \ell(t)$, $\deg(\ell) = 12$ such that $\ell(t)$ has five double zeros. It has the order 3 automorphism $\beta_\ell : (t, v) \mapsto (t, \xi v)$. The quotient $(C_\ell \times E)/(\beta_\ell \times \alpha_E)$ is a K3 surface S_ℓ which has the elliptic fibration with Weierstrass equation $Y^2 = X^3 + \ell(t)$. Then the desingularization of $(S_\ell \times E)/(\alpha_\ell \times \alpha_E)$ is a Calabi–Yau threefold X_ℓ with $h^{2,1}(X_\ell) = 4$ and $h^{1,1}(X_\ell) = 40$.

Similarly, for $k = 1$, one can find a Calabi–Yau threefold X with $h^{2,1}(X) = 5$ and $h^{1,1}(X) = 29$.

For details of these two cases, see Garbagnati–van Geemen [21] or Rohde [53].

We summarize the above discussion in the following form.

Proposition 2.32 *The Calabi–Yau threefold X_f (resp. X_ℓ) constructed above is of CM type if and only if the Jacobian variety $J(C_f)$ (resp. $J(C_\ell)$) is of CM type. In this case, X_f (resp. X_ℓ) is modular (automorphic).*

We will also mention results of Garbagnati [20], which generalize the method of Garbagnati–van Geemen [21] to automorphisms of order 4. In order to construct Calabi–Yau threefolds with a non-symplectic automorphism of order 4, start with the hyperelliptic curves

$$C_{f_g}\ :\ z^2 = tf_g(t^2),\ \deg(f_g) = g, \quad f_g \text{ without multiple roots.}$$

C_{f_g} has the automorphism $\alpha_C : (t, z) \mapsto (-r, iz)$. We consider the cases $g = 2$ or 3. Let $E_i : v^2 = u(u^2 + 1)$ be the elliptic curve and let $\alpha_E : (u, v) \mapsto (-u, iv)$ be the autmorphism of E_i. Now take the quotient of the product $E_i \times C_{f_g}/\alpha_E \times \alpha_C$. Then the singularities are of A-D-E type, and the desingularization defines a K3 surface, S_{f_g}, with the automorphism $\alpha_S : (x, y, s) \mapsto (-x, iy, s)$. In fact, there is the elliptic fibration $\mathcal{E} : y^2 = x^3 + xsf_g(s)^2$ and the map $\pi : E_i \times C_{f_g} \to \mathcal{E}$

defined by $((u,v);(z,t)) \mapsto (x := uz^2, y := vz^3, s := t^2)$ is the quotient map $E_i \times C_{f_g} \to (E_i \times C_{f_g})/\alpha_E \times \alpha_C$. The K3 surface S_{f_g} thus obtained has large Picard number. Indeed, $\mathrm{rank}(T_{S_{f_g}}) \leq 4$ if $g = 2$ and ≤ 6 if $g = 3$. Also, S_{f_g} admits the order 4 non-symplectic automorphism α_S, and the fixed loci of α_S and of α_S^2 contain no curves of genus > 0. The K3 surface (S_{f_g}, α_S^2) with involution α_S^2 indeed corresponds to the triplet $(18, 4, 1)$ for $g = 2$ and $(16, 6, 1)$ for $g = 3$.

Once we have a family of K3 surfaces with non-symplectic automorphism α_S of order 4, we can construct a family of Calabi–Yau threefolds as the quotient of the product of S_{f_g} with the elliptic curve E_i.

Proposition 2.33 *(Garbagnati [20]) There is a desingularization Y_{f_g} of $(E_i \times S_{f_g})/(\alpha_E^3 \times \alpha_S)$ which is a smooth Calabi–Yau threefold with*

$$h^{1,1}(Y_{f_g}) = \begin{cases} 73 & \text{if } g = 2 \\ 56 & \text{if } g = 3 \end{cases} \quad \text{and} \quad h^{2,1}(Y_{f_g}) = \begin{cases} 1 & \text{if } g = 2 \\ 2 & \text{if } g = 3 \end{cases}.$$

Proposition 2.34 *The Calabi–Yau threefold Y_{f_g} is of CM type if and only if the Jacobian variety $J(C_{f_g})$ is of CM type. When it is of CM type, Y_{f_g} is automorphic.*

3 The modularity of mirror maps of Calabi–Yau varieties, and mirror moonshine

• Modularity of solutions of Picard–Fuchs differential equations

Let $n \in \mathbb{N}$ and let $M_n := U_2 \oplus (-E_8)^2 \oplus < -2n >$ be a lattice of rank 19. Here U_2 is the usual hyperbolic lattice of rank 2 and $-E_8$ is the unique negative-definite unimodular lattice of rank 8, and $< -2n >$ denotes the rank 1 lattice $\mathbb{Z}v$ with its bilinear form determined by $< v, v >= -2n$. We consider a one-parameter family of M_n polarized K3 surfaces X_t over $\mathbb{Q}$ with generic Picard number $\rho(X_t) = 19$. Here by a M_n-polarized K3 surfaces, we mean K3 surfaces X_t such that $T(X_t)$ is primitively embedded into $U_2 \oplus \mathbb{Z}u$ where u is a vector of height $2n$, $n \in \mathbb{N}$. The Picard–Fuchs differential equation of X_t is of order 3. It is shown by Doran [15] that for such a family of K3 surfaces X_t, there is a family of elliptic curves E_t such that the order 3 Picard–Fuchs differential equation of X_t is the symmetric square of the order 2 differential equation associated to the family of elliptic curves E_t. The existence of such a relation stems from the so-called Shioda–Inose structures of X_t (or by Dolgachev's result [14] which asserts that the coarse moduli space of M_n-polarized K3 surfaces is isomorphic to the moduli space of elliptic curves with level n structure). Long [43] gave an algorithm how to determine a family of elliptic curves E_t, up to projective equivalence.

Yang and Yui [64] studied differential equations satisfied by modular forms of two variables associated to $\Gamma_1 \times \Gamma_2$ where Γ_i $(i = 1, 2)$ are genus zero subgroups of $SL(2, \mathbb{R})$ commensurable with $SL(2, \mathbb{Z})$. A motivation is to realize these differential equations satisfied by modular forms of two variables as Picard–Fuchs differential equations of K3 families with large Picard numbers, e.g., $19, 18, 17$ or 16, thereby establishing the modularity of solutions of Picard–Fuchs differential equations. This goal was achieved for some of the families of K3 surfaces studied by Lian and Yau in [39], [40].

• Monodromy of Picard–Fuchs differential equations of certain families of Calabi–Yau threefolds

Classically, it is known that the monodromy groups of Picard–Fuchs differential equations for families of elliptic curves and K3 surfaces are congruence subgroups of $SL(2,\mathbb{R})$. This modularity property of the monodromy groups ought to be extended to families of Calabi–Yau threefolds. For this, we will study the monodromy groups of Picard–Fuchs differential equations associated with one-parameter families of Calabi–Yau threefolds. In Chen–Yang–Yui [8] they considered 14 Picard–Fuchs differential equations of order 4 of hypergeometric type. They are of the form

$$\theta^4 - Cz(\theta+A)(\theta+1-A)(\theta+B)(\theta+10B)$$

where $A, B, C \in \mathbb{Q}$.

Theorem 3.1 *In these 14 hypergeometric cases, the matrix representations of the monodromy groups relative to the Frobenius basis can be expressed in terms of the geometric invariants of the underlying Calabi–Yau threefolds. Here the geometric invariants are the degree d, the second Chern numbers, $c_2 \cdot H$ and the Euler number, c_3.*

Furthermore, under suitable change of basis, the monodromy groups are contained in certain congruence subgroups of $Sp(4,\mathbb{Z})$ of finite index (in $Sp(4,\mathbb{Z})$) and whose levels are related only to the geometric invariants.

However, finiteness of the index of the monodromy groups themselves in $Sp(4,\mathbb{Z})$ is not established.

Using the same idea for the hypergeometric cases, the monodromy groups of the differential equations of Calabi–Yau type that have at least one conifold singularity (not of the hypergeometric type) are computed. Our calculations verify numerically that if the differential equations come from geometry, then the monodromy groups are also contained in some congruence subgroups of $Sp(4,\mathbb{Z})$.

We should mention that van Enckevort and van Straten [60] numerically determined the monodromy for 178 Calabi–Yau equations of order 4 with a different method from ours, and speculated that these equations do come from geometry.

- **Modularity of mirror maps and mirror moonshine**

For a family of elliptic curves $y^2 = x(x-1)(x-\lambda)$, the periods $\int_1^\infty \frac{dx}{\sqrt{x(x-1)(x-\lambda)}}$ satisfy the Picard–Fuchs differential equation

$$(1-\lambda)\theta^2 f - \lambda\theta f - \frac{\lambda}{4} f = 0 \quad (\theta = \lambda\frac{d}{d\lambda}).$$

The monodromy group for this Picard–Fuchs differential equation is $\Gamma(2) \subset SL(2,\mathbb{R})$ of finite index. The periods can be expressed in terms of the hypergeometric function

$${}_2F_1(\frac{1}{2},\frac{1}{2};1;\lambda).$$

Now suppose that $y_0(\lambda) = 1 + \cdots$ is the unique holomorphic solution at $\lambda = 0$ and $y_1(\lambda) = \lambda y_0(\lambda) + g(\lambda)$ be the solution with logarithmic singularity. Set $z = y_1(\lambda)/y_0(\lambda)$. Then λ, as a function of z, becomes a modular function for the modular group $\Gamma(2)$. This is called a *mirror map* of the elliptic curve family. That a mirror map is a Hauptmodul for a genus zero subgroup $\Gamma(2) \subset SL(2,\mathbb{R})$ is referred to as *mirror moonshine*.

For one-parameter families of K3 surfaces of generic Picard number 19, the Picard–Fuchs differential equations are of order 3. Since such families of K3 surfaces are equipped with Shioda–Inose structure (cf. Morrison [48]), the Picard–Fuchs differential equations are symmetric squares of differential equations of order 2. Hence the monodromy groups are realized as subgroups of $SL(2,\mathbb{R})$.

Classically, explicit period and mirror maps for the quartic K3 surface have been described by several articles, e.g., Hartmann [27] (and also see Lian and Yau [44]. Consider the deformation of the Fermat quartic: $F_t := \{x_0^4 + x_1^4 + x_2^4 + x_3^4 - 4tx_0x_1x_2x_3 = 0\} \subset \mathbb{P}^3$. Taking the quotient of F_t by some finite group and then resolving singularities, we obtain the Dwork pencil of K3 surfaces, denoted by X_t. Then X_t is a K3 surface with generic Picard number 19. The Picard–Fuchs differential equation is of order 3 and there is a unique holomorphic solution (at $t=0$) of the form $w_0(t) = 1 + \sum_{n\geq 1} c_n t^n$, and another solution of logarithmic type: $w_1(z) = \log w_0(t) + \sum_{n\geq 1} d_n t^n$. Now introduce the new variable z by $z := \frac{1}{2\pi i}\frac{w_1(t)}{w_0(t)}$, and put $q = e^{2\pi i z}$. The inverse $t = t(q)$ is the mirror map and is given by

$$t(q) = q - 104q^2 + 6444q^3 - 311744q^4 + 13018830q^5 + \cdots$$

This is the reciprocal of the Hauptmodul for $\Gamma_0(2)_+ \subset SL(2,\mathbb{R})$.

There are several more examples of one-parameter families of K3 surfaces with generic Picard nubmer 19. Doran [15] has established the modularity of the mirror map for M_n-polarized K3 surfaces. However, for each n, the explicit description of mirror maps as modular functions are still to be worked out.

The situation will get more much complicated when we consider two-parameter families of K3 surfaces. Hashimoto and Terasoma [28] have studied the period and mirror maps of the two-parameter (in fact, projective 1-parameter) family $\{\mathcal{X}_t\}$ $t = (t_0, t_1) \in \mathbb{P}^1$ of quartic family of K3 surfaces defined by

$$\mathcal{X}_t : x_1 + \cdots + x_5 = t_0(x_1^4 + \cdots + x_5^4) + t_1(x_1^2 + \cdots + x_5^2)^2 = 0$$

in $\mathbb{P}^4$ with homogeneous coordinates $(x_1 : \cdots : x_5)$. This family admits a symplectic group action by the symmetric group S_5. The Picard number of a generic fiber is equal to 19, and the Gram matrix of the transcendental lattice T is given by $\begin{pmatrix} 4 & 1 & 0 \\ 1 & 4 & 0 \\ 0 & 0 & -20 \end{pmatrix}$. The image of the period map of this family is a 1-dimensional subdomain Ω_T of the 19-dimensional period domain (the bounded symmetric domain of type IV). Let Ω_T° be a connected component of Ω_T. Since $O(T)$ has no cusp, there is a modular embedding $i : \Omega_T^\circ \to \mathbf{H}_2$ to the Siegel upper half-plane $\mathbf{H}_2$ of genus 2. This modular embedding is constructed using the Kuga-Satake construction. The inverse of the period map, that is, a mirror map, is constructed using automorphic forms of one variable on Ω_T°. In fact, automorphic forms are constructed as the pull-backs of the fourth power of theta constants of genus 2. This gives yet another example of a generalized mirror moonshine.

For a one-parameter family of Calabi–Yau threefolds, the mirror map is defined using specific solutions of the Picard–Fuchs differential equation of the family. At a point of maximal unipotent monodromy (e.g., $z=0$), there is a unique holomorphic power series solution $\omega_0(z)$ with $\omega_0(0) = 1$, and a logarithmic power series solution $\omega_1(z) = \log(z)\omega_0(z) + g(z)$ where $g(z)$ is holomorphic near $z = 0$ with $g(0) = 0$. Now put $t := \frac{\omega_1(z)}{\omega_0(z)}$. We call the map defined by $q := e^{2\pi i t} = ze^{g(z)/\omega_0(z)}$ the *mirror map* of the Calabi–Yau family. (See, for instance, Lian and Yau [44]).

For some one-parameter families of Calabi–Yau threefolds, e.g., of hypergeometric type, the integrality of the mirror maps has been established by Krattenthaler and Rivoal [35], [34].

The modularity of mirror maps of Calabi–Yau families is getting harder to deal with in general. Doran [15] has considered certain one-parameter families of Calabi–Yau threefolds with $h^{2,1}=1$. The Picard–Fuchs differential equations of these Calabi–Yau threefolds are of order 4. Under some some special constraints imposed by special geometry, and some conditions about a point $z=0$ of maximal unipotent monodromy, there is a set of fundamental solutions (to the Picard–Fuchs differential equation) of the form $\{u, u\cdot t, u\cdot F', (tF'-2F)\}$ where $u=u(z)$ is the fundamental solution locally holomorphic at $z=0$, $t=t(z)$ is the mirror map, $F(z)$ is the prepotential and F' is the derivative of F with respect to z. When there are no instanton corrections, the Picard–Fuchs differential equation becomes the symmetric cube of some second-order differential equation. In this case, Doran has shown that the mirror map becomes automorphic. However, exhibiting automorphic forms explicitly remains an open problem. On the other hand, if there are instanton corrections, a necessary and sufficient condition is presented for a mirror map to be automorphic.

We should mention here a converse approach to the modularity question of solutions of Picard–Fuchs differential equations, along the line of investigation by Yang and Yui [64]. The starting point is modular forms and the differential equations satisfied by them. It may happen that these differential equations coincide with Picard–Fuchs differential equations of some families of Calabi–Yau varieties. Consequently, the modularity of solutions of Picard–Fuchs differential equations and mirror maps can be established.

4 The modularity of generating functions of counting some quantities on Calabi–Yau varieties

Under this subtitle, topics included are enumerative geometry, Gromov–Witten invariants, and various invariants counting some mathematical/physical quantities, etc.

- Mirror symmetry for elliptic curves and quasimodular forms

We consider the generating function, $F_g(q)$, counting simply ramified covers of genus $g\geq 1$ over a fixed elliptic curve with $2g-2$ marked points.

Theorem 4.1 *For each $g\geq 2$, $F_g(q)$ (with $q=e^{2\pi i\tau}$, $\tau\in\mathfrak{H}$), is a quasimodular form of weight $6g-6$ on $\Gamma=SL_2(\mathbb{Z})$. Consequently, $F_g(q)$ is a polynomial in $\mathbb{Q}[E_2,E_4,E_6]$ of weight $6g-6$.*

This result is stated as the Fermion Theorem in Dijkgraaf [13], which is concerned with the A-model side of mirror symmetry for elliptic curves. A mathematically rigorous proof was given in the article of Roth–Yui [55]. The B-model (bosonic) counting constitutes the mirror side of the calculation. The bosonic counting will involve calculation with Feynman integrals of trivalent graphs. A mathematical rigorous treatment of the B-model counting is currently under way.

Further generalizations:

(a) The generaiting function of m-simple covers for any integer $m\geq 2$ of genus $g\geq 1$ over a fixed elliptic curve with $2g-2$ marked points has been shown again to be quasimodular forms by Ochiai [51].

(b) The quasimodularity of the Gromov–Witten invariants for the three elliptic orbifolds with simple elliptic singularities $\tilde{E}_N$ ($N = 6, 7, 8$) has been established by Milanov and Ruan [47]. These elliptic orbifolds are realized as quotients of hypersurfaces of degree $3, 4$ and 6 in weighted projective 2-spaces with weights $(1, 1, 1), (1, 2, 2)$ and $(1, 2, 3)$, respectively.

(c) The recent article of Rose [54] has proved the quasimodularity of the generating function for the number of hyperelliptic curves (up to translation) on a polarized abelian surface.

5 Future Prospects

Here we collect some topics which we are not able to cover in this paper as well some problems for further investigation.

5.1 The potential modularity. The potential modularity of families of hypersurfaces. For the Dwork families of one-parameter hypersurfaces, the potential modularity has been established by R. Taylor and his collaborators. Extend the potential modularity to more general Calabi–Yau hypersurfaces, Calabi–Yau complete intersections, etc.

5.2 The modularity of moduli of families of Calabi–Yau varieties. Moduli spaces of lattice polarized K3 surfaces with large Picard number.

5.3 Congruences, formal groups. Congruences for Calabi–Yau families, formal groups.

5.4 The Griffiths Intermediate Jacobians of Calabi–Yau threefolds.

- Explicit description of the Griffiths intermediate Jacobians and their modularity.

Let X be a Calabi–Yau threefold defined over $\mathbb{Q}$. Let

$$J^2(X) \simeq H^3(X, \mathbb{C})/F^2H^3(X, \mathbb{C}) + H^3(X, \mathbb{Z}) \simeq H^3(X, \mathbb{C})^*/H_3(X, \mathbb{Z})$$

be the Griffiths intermediate Jacobian of X. There is the Abel–Jacobi map

$$CH^2(X)_{hom,\mathbb{Q}} \to J^2(X)_{\mathbb{Q}}.$$

A part of the Beilinson–Bloch conjecture asserts that this map is injective modulo torsion.

Now suppose that X is rigid. Then $J^2(X)$ is a complex torus of dimension 1 so that there is an elliptic curve E such that $J^2(X) \simeq E(\mathbb{C})$. We know that X is modular by [26].

Question: *Is it true that the Griffiths intermediate Jacobian $J^2(X)$ of a rigid Calabi–Yau threefold X over $\mathbb{Q}$ is defined over $\mathbb{Q}$ and hence modular?*

- Special values of L-series of Calabi–Yau threefolds over $\mathbb{Q}$.

Assuming a positive answer to the above question, we can consider a possible relation between the Birch and Swinnerton-Dyer conjecture for rational points on $J^2(X)_{\mathbb{Q}}$:

$$\mathrm{rank}_{\mathbb{Z}} J^2(X)_{\mathbb{Q}}(\mathbb{Q}) = \mathrm{ord}_{s=1} L(J^2(X)_{\mathbb{Q}}, s)$$

and the Beilinson–Bloch conjecture on the Chow group $CH^2(X)_{\mathbb{Q}}$ of X:

$$\mathrm{rank}_{\mathbb{Z}} CH^2(X)_{hom,\mathbb{Q}} = \mathrm{ord}_{s=2} L(X_{\mathbb{Q}}, s).$$

If the Abel–Jacobi map $CH^2(X)_{hom,\mathbb{Q}} \to H^2(X)_{\mathbb{Q}}$ is injective modulo torsion, then

$$\mathrm{ord}_{s=2}L(X_{\mathbb{Q}}, s) \leq \mathrm{ord}_{s=1}L(J^2(X)_{\mathbb{Q}}, s).$$

5.5 Geometric realization problem (The converse problem). We know that every singular K3 surface X over $\mathbb{Q}$ is motivically modular in the sense that the transcendental cycles $T(X)$ corresponds to a newform of weight 3. For singular K3 surfaces over $\mathbb{Q}$, the converse problem asks:

Which newform of weight 3 *with integral Fourier coefficients would correspond to a singular K3 surface defined over* $\mathbb{Q}$*?*

This has been answered by Elkies and Schütt [18] (see also the article by Schütt [58] in this volume). Their result is the following theorem.

Theorem 5.1 *Every Hecke eigenform of weight* 3 *with eigenvalues in* $\mathbb{Z}$ *is associated to a singular K3 surface defined over* $\mathbb{Q}$.

Now we know that every rigid Calabi–Yau threefold over $\mathbb{Q}$ is modular (see Gouvêa–Yui [26]). The converse problem that has been raised, independently, by Mazur and van Straten is the so-called *geometric realization problem*, and is stated as follows:

Which newforms of weight 4 *on some* $\Gamma_0(N)$ *with integral Fourier coefficients would arise from rigid Calabi–Yau threefolds over* $\mathbb{Q}$*? Do all such forms arise from rigid Calabi–Yau threefolds over* $\mathbb{Q}$*?*

For Calabi–Yau threefolds, a very weak version of the above problem has been addressed in Gouvêa–Kiming–Yui [25].

Question: *Given a rigid Calabi–Yau threefold* X *over* $\mathbb{Q}$ *and a newform* f *of weight* 4*, for any non-square rational number* d*, there is a twist* f_d *by the quadratic character corresponding to the quadratic extension* $\mathbb{Q}(\sqrt{d})/\mathbb{Q}$*. Does* f_d *arise from a rigid Calabi–Yau threefold* X_d *over* $\mathbb{Q}$*?*

A result of Gouvêa–Kiming–Yui [25] in this volume is that the answer is positive if a Calabi–Yau threefold has an anti-symplectic involution. Let X be a rigid Calabi–Yau threefold over $\mathbb{Q}$. For a square-free $d \in \mathbb{Q}^{\times}$, let $K := \mathbb{Q}(\sqrt{d})$ and let σ be the non-trivial automorphism of K. We say that a rigid Calabi–Yau threefold X_d defined over $\mathbb{Q}$ is a *twist of* X *by* d if there is an involution ι of X which acts by -1 on $H^3_{et}(\bar{X}, \mathbb{Q}_\ell)$, and an isomorphism $\theta : (X_d)_K \cong X_K$ defined over K such that $\theta^\sigma \circ \theta^{-1} = \iota$.

Proposition 5.2 *Let* X *be a rigid Calabi–Yau threefold over* $\mathbb{Q}$ *and let* f *be the newform of weight* 4 *attached to* X*. Then, if* X_d *is twist by* d *of* X*, the newform attached to* X_d *is* f_d*, the twist of* d *by the Dirichlet character* χ *corresponding to* K*.*

Various types of modular forms have appeared in the physics literature. We wish to understand "conceptually" why modular forms play such pivotal roles in physics. Here we list some of the modular appearances in the physics literature.

5.6 Modular forms and Gromov–Witten invariants. Modular (automorphic) forms and Gromov–Witten invariants and generalized invariants.

5.7 Automorphic black hole entropy. This has something to do with Conformal Field Theory. Mathematically, mock modular forms, Jacobi forms etc. will come into the picture. This is beyond the scope of this article.

5.8 M_{24}**-moonshine.** Recently, some close relations between the elliptic genus of K3 surfaces and the Mathieu group M_{24} along the line of moonshine have been observed in the physics literature, e.g., [16],[17]. It has been observed that multiplicities of the non-BPS representations are given by the sum of dimensions of irreducible representations of M_{24} and furthermore, they coincide with Fourier coefficients of a certain mock theta function.

6 Appendix

In this appendix, we will recall modular forms of various kinds, e.g., classical modular forms, quasimodular forms, Hilbert modular forms, Siegel modular forms, and most generally, automorphic forms, which are relevant to our discussions. For details, the reader is referred to [5].

Definition 6.1 Dimension 1: Let $\mathfrak{H} := \{z \in \mathbb{C} \,|\, \mathrm{Im}(z) > 0\}$ be the complex upper-half plane. For a given integer $N > 0$, let

$$\Gamma_0(N) =: \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,\mathbb{Z}) \,|\, c \equiv 0 \mod N \right\}$$

be a congruence subgroup of $SL(2,\mathbb{Z})$ (of finite index). A modular form of *weight* k and *level* N is a holomorphic function $f : \mathfrak{H} \to \mathbb{C}$ with the following properties:

(M1) For $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N)$, $f\left(\frac{az+b}{cz+d}\right) = (cz+d)^k f(z)$;

(M2) f is holomorphic at the cusps.

Since $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \in \Gamma_0(N)$, (a1) implies that $f(z+1) = f(z)$, so f has the Fourier expansion $f(z) = f(q) = \sum_{n\geq 0} c(n)q^n$ with $q = e^{2\pi i z}$. f is a *cusp* form if it vanishes at all cusps.

If χ is a mod N Dirichlet character, we can define a modular form with character χ by replacing (a1) by $f\left(\frac{az+b}{cz+d}\right) = \chi(d)(cz+d)^k f(z)$.

The space $S_k(\Gamma_0(N))$ of all cusp forms of weight k and level N is a finite dimensional vector space, and similarly, so also is the space $M_k(\Gamma_0(N),\chi)$ of all modular forms of weight k and level N.

On $S_k(\Gamma_0(N),\chi)$ there are Hecke operators T_p for every prime p not dividing N. A cusp form f is a (normalized) Hecke *eigenform* if it is an eigenvector for all T_p, that is, $T_p(f) = c(p)f$. For such a normalized eigenform f, define the L-series $L(f,s)$ by

$$L(f,s) = \sum_{n\geq 1} c(n)n^{-s} = \prod_p \frac{1}{1 - c(p)p^{-s} + \chi(p)p^{k-1-2s}} \quad \text{where } \chi(p) = 0 \text{ if } p|N$$

Dimension 2: Let $F = \mathbb{Q}(\sqrt{d})$ be a totally quadratic field over $\mathbb{Q}$ where $d > 0$ and square-free, and let $\mathcal{O}_F$ be its ring of integers. The $SL_2(\mathcal{O}_F)$ can be embedded into $SL_2(\mathbb{R}) \times SL_2(\mathbb{R})$ via the two real embeddings of F to $\mathbb{R}$, and it acts on $\mathfrak{H} \times \mathfrak{H}$ via fractional linear transformations:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} z = \left(\frac{az_1 + b}{cz_1 + d}, \frac{az_2 + b}{cz_2 + d} \right)$$

for $z = (z_1, z_2) \in \mathfrak{H} \times \mathfrak{H}$. The group

$$\Gamma(\mathcal{O}_F \oplus \mathfrak{a}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(F), a, d \in \mathcal{O}_F, b\mathfrak{a}^{-1}, c \in \mathfrak{a} \right\}$$

is called *the Hilbert modular group* corresponding to a fractional ideal $\mathfrak{a}$ of F. If $\mathfrak{a} = \mathcal{O}_F$, put $\Gamma_F = \Gamma(\mathcal{O}_F \oplus \mathcal{O}_F) = SL_2(\mathcal{O}_F)$. Let $\Gamma \subset SL_2(F)$ be a subgroup commensurable with Γ_F, and let $(k_1, k_2) \in \mathbb{Z} \times \mathbb{Z}$.

A meromorphic function $f : \mathfrak{H} \times \mathfrak{H} \to \mathbb{C}$ is called a meromorphic *Hilbert modular form* of *weight* (k_1, k_2) for Γ if

$$f(\gamma\, z) = (cz_1 + d)^{k_1} (cz_2 + d)^{k_2} f(z)$$

for all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ and $z = (z_1, z_2) \in \mathfrak{H}^2$. If f is holomorphic, then f is a holomorphic Hilbert modular form, and a holomorphic Hilbert modular form is *symmetric* if $f(z_1, z_2) = f(z_2, z_1)$. Further, f is a cusp form if it vanishes at cusps of Γ. A cusp form of weight $(2, 2)$ is identified with a holomorphic 2-form on the Hilbert modular surface $\mathfrak{H}^2/\Gamma$. The space of holomoprhic Hilbert modular forms of weight (k_1, k_2) for Γ is denoted by $M_k(\Gamma)$ and the cusp forms by $S_k(\Gamma)$. $M_k(\Gamma)$ is a finite dimensional vector space over $\mathbb{C}$.

A holomorphic Hilbert modular form has a Fourier expansion at ∞ of the following form. Let $M \subset F$ be a rank 2 lattice and $V \subset \mathcal{O}_F^*$ be a finite index subgroup acting on M in a suitable way. Then

$$f(z) = a_0 + \sum_{\nu \in M^\vee,\, \nu \geq 0} a_\nu e^{2\pi i \mathrm{tr}(\nu z)}$$

where ν runs over the dual lattice $M^\vee$, and $\mathrm{tr}(\nu\, z) := v z_1 + v' z_2$.

The L-series of a Hilbert modular cusp form is defined by

$$L(f, s) = \sum_{\mathfrak{a} \subset \mathcal{O}_F} a(\mathfrak{a}) N(\mathfrak{a})^{-s}$$

where $\mathfrak{a}$ runs over principal ideals. For details, the reader should consult Bruinier [6].

Dimension 3: Let $g, N \in \mathbb{N}$. Define the Siegel upper-half plane by

$$\mathfrak{H}_g := \left\{ z \in M_{g \times g}(\mathbb{C}) \,|\, z^t = z, \mathrm{Im}(z > 0 \right\},$$

and the symplectic group $Sp(2g, \mathbb{Z})$ as the automorphism group of the symplectic lattice $\mathbb{Z}^{2g}$. That is,

$$Sp(2g, \mathbb{Z}) = \{ \gamma \in GL(2g, \mathbb{Z}) \,|\, \gamma^t J \gamma = J \}$$

where $J_g := \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix}$. The group $Sp(2g, \mathbb{Z})$ acts on $\mathfrak{H}_g$ by

$$z \mapsto \gamma(z) = (Az + B)(Cz + D)^{-1} \quad \text{for} \quad \gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(2g, \mathbb{Z}).$$

Let $\Gamma_g(N) := \{ \gamma \in Sp(2g, \mathbb{Z}) \,|\, \gamma \equiv I_{2g} \pmod N \}$ be a subgroup of $Sp(2g, \mathbb{Z})$. Then $\Gamma_g(N)$ acts freely when $N \geq 3$. Here I_{2g} is the identity matrix of order $2g$. If

$N = 1, \Gamma_g(1) = Sp(2g, \mathbb{Z})$. The quotient space $\mathfrak{H}_g/\Gamma_g(N)$ (with $N \geq 3$) is a complex manifold of dimension $g(g+1)/2$ (associated to a graded algebra of modular forms).

A holomorphic function $f : \mathfrak{H}^g \to \mathbb{C}$ is a *Siegel modular form of genus g, weight* $k \in \mathbb{N}$ if

$$f(\gamma(z)) = \det(Cz + D)^k f(z)$$

for all $\gamma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(2g, \mathbb{Z})$ an d all $z \in \mathfrak{H}_g$. The space of all holomorphic Siegel modular forms is a finitely generated graded algebra. The simplest examples of Siegel modular forms are given by theta constants.

A holomorphic Siegel modular form f has a Fourier expansion of the form

$$f(z) = \sum A(n) e^{2\pi \mathrm{tr}(nz)}$$

where the sum runs over all positive semi-definite integral matrices $n \in GL(g, \mathbb{Q})$. If the Fourier expansion is supported only on positive definite integral $g \times g$ matrices n, then f is called a cusp form.

There are at least two different L-series of a holomorphic Siegel modular form f: one is the spinor L-series, and the other is the standard L-series.

However, it is not clear how to associate these L-series to the Fourier expansion.

For details, the reader should consult van der Geer [22].

Example 6.2 (a) For $N = 1$, the total space $M := \oplus_k M_k(SL(2, \mathbb{Z}))$, of all modular forms is generated by the Eisenstein series E_4 and E_6. The Eisenstein series E_2 is not modular but it may be called *quasimodular*. The space $\tilde{M}$ of all quasimodular forms for $SL(2, \mathbb{Z})$ is generated by the Eisenstein series E_2, E_4 and E_6, that is, $\tilde{M} = \mathbb{C}[E_2, E_4, E_6]$.

For $N > 1$, the space of modular forms of weight k for any congruence subgroup of level N is finite dimensional, and its basis can be determined.

Some properties of quasimodular forms for finite index subgroups of $SL_2(\mathbb{Z})$, dimension, basis, etc.

(b) There are Eisenstein series for $\Gamma_F = SL_2(\mathcal{O}_F)$ and k even given by

$$G_{k,B} = N(\mathfrak{b})^k \sum_{(c,d) \in \mathcal{O}_F^* \backslash \mathfrak{b}^2} N(cz + d)^{-k}$$

for B an ideal class of F.

If we put $g_k := \frac{1}{\zeta_F(k)} G_{k,\mathcal{O}_F}$, then g_2, g_6 and g_{10} generate the graded algebra $M_{2^*}^{symm}(\Gamma_F)$ over $\mathbb{C}$, that is,

$$M_{2^*}^{symm}(\Gamma_F) \cong \mathbb{C}[g_2, g_6, g_{10}].$$

(c) (Igusa) For $g = 2$, the graded algebra $\mathfrak{M}$ of classical Siegel modular forms of genus 2 is generated by the Eisenstein series E_4 and E_6, the Igusa cusp forms C_{10}, C_{12}, and C_{35} (where the subindex denote weights), and

$$\mathfrak{M} \cong \mathbb{C}[E_4, E_6, C_{10}, C12, C_{35}]/(C_{35}^2 = P(E_4, E_6, C_{10}, C_{12}))$$

where P is an explicit polynomial.

Acknowledgments

I would like to thank Matthias Schütt for carefully reading the preliminary version of this paper and suggesting numerous improvements. I am indebted to Ron Livné for answering my questions about Galois representations.

I would also like to thank a number of colleagues for their comments and suggestions. This includes Jeng-Daw Yu, Ling Long, Ken-Ichiro Kimura, and Bert van Geemen.

We are grateful to V. Nikulin for allowing us to use the template of Nikulin's pryamid in Figure 1.

Last but not least, my sincere thanks is to the referee for reading through the earlier versions of this article and for very helpful constructive criticism and suggestions for the improvement of the article. I would also like to thank Arther Greenspoon of Mathematical Reviews for copy-editing the article.

The article was completed while the author held visiting professorship at various institutions in Japan: Tsuda College, Kavli Institute for Physics and Mathematics of the Universe and Nagoya University. I thank the hospitality of these instutions.

References

[1] Artebani, M., Boissière, S. and Sarti, A. *The Berglund–Hübsch–Chiodo–Ruan mirror symmetry for K3 surfaces*, arXiv:1108.2780.

[2] Bini, C., and van Geemen, B. *Geometry and arithmetic of Maschke's Calabi–Yau threefold*, Commun. Number Theory Phys. **5**, No.4 (2011).

[3] Borcea, C. *Calabi–Yau threefolds and complex multiplication*, in *Essays on Mirror Manifolds*, pp. 489–502. Int. Press, 1992.

[4] Borcea, C. *K3 surfaces with involution and mirror pairs of Calabi–Yau manifolds*, in *Mirror Symmetry*, AMS/IP Stud. Adv. Math. **1**, Amer. Math. Soc. 1997, pp. 717–743. **33** (1983), 227–250.

[5] Bruiner, J., van der Geer, G., Harder, G., and Zagier, D., *The 1-2-3 of Modular Forms*, Universitext Springer-Verlag, Berlin 2008.

[6] Bruinier, J. *Hilbert modular forms and their applications*, in *The 1-2-3 of Modular Forms*, Universitext. Springer, Berlin 2008, pp. 105–179.

[7] Consani, C. and Scholten, J. *Arithmetic on a quintic threefold*, Internat. J. Math. **12** (8) (2001), pp. 943–972.

[8] Chen Y.-H., Yang, Y. and Yui, N., *Monodromy of Picard–Fuchs differential equations for Calabi–Yau threefolds*, J. Reine Angew Math. **616** (2008), 167–203.

[9] Cynk, S., and Meyer, C. *Modularity of some nonrigid double octic Calabi–Yau threefolds*, Rocky Mountain J. Math. **38** (2008), pp. 1937–1958.

[10] Deligne, P., Milne, J.S., Ogus, A., and Shih, K.-Y. *Hodge cycles, motives and Shimura varieties*, Lecture Notes in Math, **900**, Springer-Verlag 1982.

[11] Dieulefait, L. *On the modularity of rigid Calabi–Yau threefolds*, Zap. Nauchn. Sem. S-Peterburg. Otdel, Mat. Inst. Steklov. (POMI) **377** (2010), Issledovaniya po teorii Chisel. 10, pp. 44–49.

[12] Dieulefait, L., Pacetti, A. and Schütt, M. *Modularity of the Consani–Scholten quintic*, arXiv:1005.4523, 2010.

[13] Dijkgraaf, R., *Mirror symmetry and elliptic curves*, in *The moduli space of curves* (Texel Island, 1994), Prog. Math. **129**, 1995. pp. 149–163.

[14] Dolgachev, I. *Mirror symmetry for lattice polarized K3 surfaces. Algebraic Geometry, 4*, J. Math. Sci **81** (1996), No. 3, pp. 2599–2630.

[15] Doran, C. *Picard–Fuchs uniformization: modularity of the mirror map and mirror-moonshine*, in *The arithmetic and geometry of algebraic cycles* (Banff AB, 1998), pp. 257–281. CRM Proc. Lecture Notes **24**, Amer. Math. Soc. 2000.

[16] Eguchi, T., and Hikami, K. *Superconformal algebras and mock theta functions*, J. Phys. A. **42**, no. 30, 304010 (2009).

[17] Eguchi, T., and Hikami. K. *Superconformal algebras and mock theta functions 2. Rademacher expansion for K3 surface*, Commun. Number Theory Phys **3** (2009), pp. 531–554.

[18] Elkies, N., and Schütt, M. *Modular forms and K3 surfaces*, arXiv:0809.0830, 2008.

[19] Freitag, E. and Salvati Manni, R. *Some Siegel threefolds with a Calabi–Yau model*, Ann Sc. Norm Super. Pisa Cl. Sci. **9** (2010), pp. 833–850, *On Siegel threefolds with a projective Calabi–Yau model*, Commun. Number Theory Phys. **5** (2011), pp. 713–750.

[20] Garbagnati, A. *New examples of Calabi–Yau* 3*-folds without maximal unipotent monodromy*, arXiv:1005.0094. May 2010.

[21] Garbagnati, A., and van Geemen, B. *Examples of Calabi–Yau threefolds parametrized bu Shimura varieties*, Rend. Sem. Mat. Univ. Pol. Torino **68** (2010), pp. 271–287.

[22] van der Geer, G. *Siegel modular forms and their applications*, in *The 1-2-3 of Modular Forms*, Universitext, Springer, 2008, pp. 181–245.

[23] Goto, Y., Livné, R., and Yui, N. *The modularity of certain K3 fibered Calabi–Yau threefolds over* $\mathbb{Q}$ *with involution*, preprint 2011.

[24] van Geemen, B. and Nygaard, N. *On the geometry and arithmetic of some Siegel modular threefolds*, J. Number Theory **53** (1995), pp. 45–87.

[25] Gouvêa, F., Kiming, I., and Yui, N., *Quadratic twists of rigid Calabi–Yau threefolds over* $\mathbb{Q}$, arXiv: 1111.5275. In this volume.

[26] Gouvêa, F. and Yui, N. *Rigid Calabi–Yau threefolds over* $\mathbb{Q}$ *are modular*, Expos. Math. **29** (2011), pp. 142–149.

[27] Hartmann, H., *Period-and mirror maps for the quartic K3*, arXiv:1101.4601, 2011.

[28] Hashimoto, K., and Terasoma, T., *Period maps of a certain K3 family with an* S_5*-action*, J. Reine Angew Math. **652** (2011), pp. 1–65.

[29] Hulek, K., and Verrill, H. *On the modularity of Calabi–Yau threefolds containing elliptic ruled surfaces*, in *Mirror Symmetry V*, AMS/IP Stud. Adv. Math. **38**, Amer. Math. Soc. 2006, pp. 19–34.

[30] Hulek, K., and Verrill, H. *On modularity of rigid and non-rigid Calabi–Yau varieties associated to the root lattice* A_4, Nagoya J. Math. **179** (2005), pp. 103–146.

[31] Khare,C. and Wintenberger, J.-P. *Serre's modularity conjecture, I and II*, Invent. Math. **178** (2009), pp. 485–504, and pp. 505–586.

[32] Kisin, M. *Modularity of* 2*-adic Barsotti–Tate representations*, Invent. Math. **178** (2009), pp. 587–634.

[33] Kondo, S. *Automorphisms of algebraic K3 surfaces which act trivially on Picard groups*, J. Math. Soc. Japan **44** (1992), pp. 75–98.

[34] Krattenthaler, C., and Rivoal, T. *On the integrality of the Taylor coefficients of mirror maps, I*, Duke J. Math. **151**, Np. 2 (2010), pp. 175–218.

[35] Krattenthaler, C., and Rivoal, T. *On the integrality of the Taylor coefficients of mirror maps, II*, Commun. Number Theory and Phys. **3**, No.3 (2009), pp. 555–591.

[36] Lee, E. *A modular non-rigid Calabi–Yau threefold*, in *Mirror Symmetry V*, AMS/IP Stud. Adv. Math. **38**, Amer. Math. Soc. 2006, pp. 89–122.

[37] Lee, E. *Update on modular non-rigid Calabi–Yau threefolds*, in *Modular Forms and String Duality*, Fields Institute Communications Vol. $\underline{5}$4, pp. 65–81.

[38] Lee, E. *A modular quintic Calabi–Yau threefold of level* 55, Canad. J. Math. **63** (2011), pp. 616–633.

[39] Lian, B.-H. and Yau, S.-T. *Mirror maps, modular relations and hypergeometric series, I*, in *XIth International Congress of Mathematical Physics (Paris 1994)*, Int. Press 1995. pp. 163–184.

[40] Lian, B.-H. and Yau, S.-T. *Mirror maps, modular relations and hypergeometric series, II*, NUclear Phys. B Proc. Suppl. **46** (1996), pp. 248–262.

[41] Livńe, R. *Motivic orthogonal two-dimensional representations of* $Gal(\bar{\mathbb{Q}}/\mathbb{Q})$, Israel J. Math. **92** (1995), pp. 149–156.

[42] Livné, R., Schuett, M., and Yui, N. *The modularity of K3 surfaces with non-symplectic group action*, Math. Annalen **348** (2010), pp.333–355.

[43] Long, L. *On Shioda–Inose structures of one-parameter families of K3 surfaces*, J. Number Theory **109** (2004), pp. 299–318.

[44] Lian, B.-H, and Yau, S-T., *Arithmetic properties of mirror maps and quantum coupling*, Comm. Math. Phys. **176** (1) (1996), pp. 163–191.

[45] Ma, S. *Rationality of the moduli spaces of* 2*-elementary K3 surfaces*, arXiv:11105.5110.

[46] Meyer, C. *Modular Calabi–Yau threefolds*, Fields Institute Monograph **22**, 2005, American Math. Soc.

[47] Milanov, T., and Ruan, Yongbin. *Gromov–Witten theory of elliptic orbifold* $\mathbb{P}^1$ *and quasi-modular forms*, arXiv:1106.2321.

[48] Morrison, D. *On K3 surfaces with large Picard number*, Invent. Math. **75** (1984), pp. 105–121.

[49] Nikulin, V. *Integral symmetric bilinear forms and some of their geometric applications*, Math. USSR-Izv. **14** (1980), pp. 103–167.

[50] Nikulin, V. *Discrete reflection groups in Lobachesvsky spaces and algebraic surfaces*, Proc. ICM Berkeley, California 1986, pp. 654–671.

[51] Ochiai, H. *Counting functions for branched covers of elliptic curves and quasimodular forms*, RIMS Kokyuroko **1218** (2001), pp. 153–167.

[52] Reid, M. *Cananical 3-folds*, Proc. Alg. Geom. Anger 1979, Sijthoff and Nordhoff, pp. 273–310.

[53] Rohde, C. *Cyclic coverings, Calabi–Yau manifolds and complex multiplication*, Lecture Notes in Math. **1975**, Springer 2009.

[54] Rose, S. *Counting hyperelliptic curves on an abelian surface with quasi-modular forms*, arXiv:1202.2094.

[55] Roth, M., and Yui, N., *Mirror symmetry for elliptic curves: The A-model (Fermionic) counting*, Clay Math. Proc. **12** (2010), pp. 245–283.

[56] Schütt, M. *On the modularity of three Calabi–Yau threefolds with bad reduction at* 11, Canad. Math. Bull. **49** (2006), pp. 296–312,

[57] Schütt, M. *Modularity of Maschke's octic and Calabi–Yau threefold*, Commun. Number Theory Phys. **5**, No.4 (2011), pp. 827–848.

[58] Schütt, M. *Two lectures on the arithmetic of K3 surfaces*, in this volume.

[59] Tate, J. *Conjectures on algebraic cycles in* ℓ*-adic cohomology*, in Motives, Proceedings Symp. Pure Math. **55**, Part I, pp. 71–83. Amer. Math. Soc. 1994.

[60] van Enckevort, C., and van Straten, D. *Monodromy calculations of fourth order equations of Calabi–Yau type*, in *Mirror Symmetry V*, Proceedings of BIRS workshop on Calabi–Yau Varieties and Mirror Symmetry, December 6–11, 2003. AMS/IP Stud. Adv. Math. **38** (2006), pp. 530–550.

[61] Voisin, C. *Mirrors et involutions sur les surfaces K3*, Astérisque **218** (1993), pp. 273–323.

[62] Vorontsov, S.P. *Automorphisms of even lattices arising in connection with automorphisms of algebraic K3 surfaces (Russian)*, Vestnik Moskov. Uni. Ser. I mat. Mekh. No.2 (1983), pp. 19–21.

[63] Yonemura, H. *Hypersurface simple K3 singularities*, Tôhoku J. Math. **42** (1990), pp. 351–380.

[64] Yang, Y., and Yui, N., *Differential equations satisfied with modular forms and K3 surfaces*, Illinois J. Math. **51** (2) (2007), pp. 667–696.

[65] Yui, N. *Update on the modularity of Calabi–Yau varieties*, in *Calabi–Yau Varieties and Mirror Symmetry*, Fields Institute Communications Vol. **38** (2003), pp. 307–362.

[66] Zarhin, Yu.G. *Hodge groups of K3 surfaces*, J. Reine Angew. Math. **341** (1983), pp. 193–220.